

DIRECTIONS FOR USE

Crops	Weeds Controlled	State	Rate kg / ha	Critical Comments
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed,	All States	4.5 kg in 330 – 550L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.
Transplanted	Portulaca, Winter grass, Rye grass, Summer Grass, Wireweed, Prince of Wales feather, Shepherd's Purse, Nettles.	Vic only	2.3 kg in 300 – 550L	Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours. Apply on light sandy soils, low in organic matter
Turf & lawns Common couch	Winter grass	All States	1.2 kg (12g/100m ²)	Rate/Bowling Green 37 m x 37 m 170g
Queensland blue couch		Qld only	0.6 kg (6g/100m ²)	85g
Buffalo grass		WA only	1.2 kg (12g/100m ²)	170g
Lucerne, Clover, Medics etc. Grown for forage, Hay or Seed production	Annual Ryegrass, Winter Grass, Barley Grass, Silver grass. Prairie Grass, Canary Grass, Great Brome, Squirrel tail Fescue and Rats tail Fescue. Wild oats, Perennial Ryegrass, Bent grass, English couch, Yorkshire fog grass, Chickweed, Seedling Sorrel, Blackberry nightshade, Patterson's curse, seedling Wireweed.	NSW, ACT, S A, W A, Tas only	1 – 1.5 kg	SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July. WEEDS: Apply when weeds are at post or Pre-emergent stages. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.
			1.5 – 2 kg	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

WEED RESISTANCE WARNING

GROUP K HERBICIDE

FMC Propyzamide WG Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is a Group K herbicide. Some naturally occurring weed biotypes resistant to FMC Propyzamide WG Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FMC Propyzamide WG Herbicide or other amide herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, FMC Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of FMC Propyzamide WG Herbicide to control the resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a FMC Crop Protection Pty Ltd representative.

Spray Preparation

FMC Propyzamide WG Herbicide is a water dispersible granule formulation and must be thoroughly dispersed in water and applied as a dilute spray solution.

- Partly fill the boomspray-tank with water so to form a slurry.
- Add the required amount of FMC Propyzamide WG Herbicide for the area to be sprayed into water containers.
- Add this correct amount of pre-mixed FMC Propyzamide WG Herbicide into the boomspray-tank with the agitation system operating.
- Fill up to the correct volume with water. The mixture must be kept in suspension at all times by continuous agitation, otherwise bottom settling of the product will result. Re-dispersion of the product will be extremely difficult and blockages of filters are likely to occur.
- When mixing with other chemical products ensure to check all label instructions prior to chemicals being tank mixed with FMC Propyzamide WG Herbicide to check for chemical compatibility. Test a small area for any adverse affects prior to application.

DO NOT mix with any products that contains the active constituent, MCPA. If tank mixing, FMC Propyzamide WG Herbicide must be in suspension before adding other herbicides, insecticides or surfactant. Check all labels Directions for Use table prior to tank mixing with Rainbow Propyzamide 500 WG Herbicide. Wash sprayer thoroughly after use.

Equipment

Always ensure the sprayer is clean, using a suitable spray and boom cleaner as directed.

It is recommended that for turf situations, up to 100 L/ha spray volume is required.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply to, or allow spray to drift onto crops, pastures or non target – desirable plants, nor onto agricultural land scheduled for cropping or pasture, as injury to the crop or pasture may occur.

PROTECTION OF LIVESTOCK

DO NOT graze sheep, cut for stock feed (including poultry and pigs) any treated areas (for 25 days after application).

DO NOT FEED TURF / LAWN CLIPPINGS TO POULTRY OR LIVESTOCK

Storage and Disposal

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. **DO NOT** re-use container or dispose of undiluted chemicals on-site. **DO NOT** dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation, and tree roots. Empty containers and product should not be burnt.

Safety Directions

Will irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling the dust. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

MSDS

For further information on FMC Propyzamide WG Herbicide refer to the Material Safety Data Sheet (MSDS) which is available from the supplier or from fmccrop.com.au.

WARRANTY

FMC Crop Protection Pty Ltd makes no warranty expressed or implied, concerning the use of this product other than that indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

APVMA Approval Number: 68080/57451

© FMC is a Registered Trademark of FMC Corporation Pty Ltd.



FMC Crop Protection Pty Ltd.
A.B.N. 48 159 288 123
5 Palmer Place
Murarrie QLD 4172
Phone: 1800 066 355
www.fmccrop.com.au
Technical Enquiries: 1800 901 939

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA WIDE
---	--

BN:

DOM: