



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

ACTIVE CONSTITUENT:

69 g/L Fenoxaprop-P-Ethyl, 34.5 g/L Cloquintocet-Mexyl

PACK SIZES:

10 – 1000 L

SOLVENT: 390 g/L Liquid Hydrocarbons



GROUP	A	HERBICIDE
-------	---	-----------

For the selective control of wild oats and annual phalaris in wheat, barley, triticale and cereal rye crops as specified in the Directions for Use Table.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

If swallowed, **DO NOT** induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet that can be obtained from www.fmccrop.com.au

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

GENERAL INSTRUCTIONS

Results are best under good growing conditions. Application to weeds or crop under stress (eg. due to continuous severe frosts, nutrient deficient, diseased, dry or waterlogged conditions) should be avoided. Generally the best yield advantage will come from controlling weeds as early as possible.

RESISTANT WEEDS WARNING

GROUP	A	HERBICIDE
-------	---	-----------

FOXTROT® Selective Herbicide is a member of the aryloxyphenoxypropionate group of herbicides. FOXTROT® is a herbicide which inhibits production of acetyl CoA carboxylase. For weed resistance management FOXTROT® is a Group A herbicide. Some naturally occurring weed biotypes resistant to FOXTROT®, and other Group A herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FOXTROT® or other Group A herbicides. Since occurrence of resistant weeds is difficult to detect prior to use, FMC accepts no liability for any losses that may result from the failure of FOXTROT® to control resistant weeds. **DO NOT** make more than one application of a Group A Herbicide to a crop per

season. For further information on herbicide resistance management consult FMC representative, trained agricultural consultant or local department agronomist.

EXPORT OF TREATED PRODUCE

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with FOXTROT® Selective Herbicide. If you are growing produce for export, please check with FMC for the latest information on MRLs and import tolerances BEFORE using FOXTROT® Selective Herbicide.

APPLICATION

Aim for complete coverage of weeds with FOXTROT®. Follow directions under "Equipment".

MIXING

Add the required quantity of FOXTROT® directly to the spray tank when about two-thirds full. Top up with water to the required volume, with agitators in motion, after adding broadleaf herbicides if required.

EQUIPMENT

Ground sprayers - Standard boom sprays only are recommended and must be fitted with by-pass or mechanical agitation. It is recommended to use approximately 50 to 100 L of water/ha and a droplet size of 200 to 300 microns. Do not apply with boomless jets or misters.

Aircraft – FOXTROT® may be applied by aircraft. Best results are obtained using 25 to 30 L of water per hectare with a steady cross wind. Do not exaggerate the swath width. Aim for a droplet size of 200 to 300 microns.

COMPATIBILITY

Mixtures of FOXTROT® with broadleaf herbicides may lead to a reduction in grass weed control. This reduction can be minimised by closely following all use recommendations and restrictions.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to the aquatic environment. **DO NOT** contaminate streams, rivers or waterways with this product or the used container.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply in weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT plant susceptible crops (eg. corn, sorghum, oats or barley) for 3 weeks after application.

Undersown Clovers and Medics – FOXTROT® does not affect undersown clovers or medics or other broadleaf crops.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour. When preparing spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat and elbow-length PVC gloves. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

DIRECTIONS FOR USE

RESTRAINT:

DO NOT apply if rainfall is expected within 4 hours.

DO NOT apply to weeds or crop under stress due to, for example, very dry, very wet, nutrient deficient, frosty or diseased conditions.

CROP	WEED	RATE (mL/ha)	CRITICAL COMMENTS
Wheat, Barley, Triticale, Cereal Rye	Wild oats (<i>Avena spp.</i>)	475	Apply 3 to 4 weeks after sowing when cereal has 2 to 5 true leaves (Z12 to Z15). Do not use this rate when wild oats are past the four leaf stage (Z14) or when there are more than 200 plants per square metre.
		560	Apply generally 3 to 6 weeks after sowing when cereal is usually 2 leaf to early tillering (Z12 to Z22). Aim to apply to actively growing young weeds up to 5 total leaves (1 tiller) (Z14, Z21).
		635	Apply up to approximately 10 weeks after sowing to actively growing weeds up to mid-tillering (Z22 to Z24)
	Annual phalaris (<i>Phalaris minor</i>) Paradoxa grass (<i>Phalaris paradoxa</i>)	635	Apply generally 3 to 4 weeks after sowing when cereal usually has 2 to 5 true leaves (Z12 to Z15). Apply only to weed densities below 200 plants per square metre. Aim to apply to actively growing young weeds up to 4 total leaves (Z14).
		800	Apply generally 4 to 6 weeks after sowing when cereal has usually reached early tillering (Z21 to Z22). Aim to apply to actively growing young weeds up to 5 total leaves (1 tiller) (Z14, Z21).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Harvest: DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION

Grazing: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 WEEKS AFTER

NOTICE TO BUYER

To the extent permitted by the Competition and Consumer Act (2010) or any relevant legislation of any State or Territory (the "Legislation") all conditions and warranties and statutory or other rights of action, whether arising in contract or tort or whether due to the negligence of FMC or Seller, which buyer or any other user may have against FMC or Seller are hereby excluded provided however that any rights of the buyer pursuant to non-excludable conditions or warranties of the Legislation are expressly preserved. FMC hereby gives notice to buyer and other users that to the extent permitted by the Legislation it will not accept responsibility for any indirect or consequential loss of whatsoever nature arising from the storage, handling or use of this Product. Where permitted by the Legislation FMC's liability shall in all circumstances be limited to the replacement of the product, or a refund of the purchase price paid therefor.

The Product must be used and applied strictly in accordance with the label instructions and other directions for use. It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of FMC or the Seller. Buyer accepts these risks.

IN A MEDICAL EMERGENCY CALL 1800 033 111 ALL HOURS

The information set forth herein is furnished free of charge and is based on technical data that FMC believes to be reliable. It is intended for use by persons having technical skill at their own discretion and risk.

FMC makes no warranties, expressed or implied, and assumes no liability in connection with any of this information. Nothing herein is to be taken as license to operate under or a recommendation to infringe on any patents.

Copyright© FMC Corporation. All rights reserved.

The FMC logo and Foxtrot® are Trademarks or Registered trademarks of FMC or its affiliates.

* Non FMC Trademark

APVMA Approval No: 61410/0607

FMC Australasia Pty Ltd.
Level 2, Building B
12 Julius Ave
North Ryde NSW 2113
Phone: 1800 066 355
www.fmccrop.com.au

SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA WIDE
