

POISON**KEEP OUT OF REACH OF CHILDREN****READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Vincit[®] C

Seed Dressing Flowable

**ACTIVE CONSTITUENTS: 25 g/L FLUTRIAFOL
4 g/L CYPERMETHRIN**

GROUP**3****FUNGICIDE****GROUP****3A****INSECTICIDE**

**Seed dressing for treatment of barley, oats, wheat and triticale
as per Directions for Use**

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT

Contents 10L

Cheminova Australia Pty Ltd ABN 59 472 616 773 / ACN 110 199 169
12 Julius Avenue, North Ryde, NSW 2113
Telephone: 1800 624 597 • www.cheminova.com.au



140/430-10F

Batch No:

Date of Manufacture

 **CHEMINOVA**
HELPING YOU GROW

DIRECTIONS FOR USE

RESTRAINT: Do NOT apply 'Vydate' to seed pretreated with this product.

CROP	DISEASE CONTROLLED	STATE	RATE/ 100 kg SEED	CRITICAL COMMENTS
Wheat	Bunt (<i>Tilletia foetida</i> , <i>T. caries</i>), Loose smut (<i>Ustilago segetum</i> var <i>tritici</i>), Seedborne flag smut (<i>Urocystis agropyri</i>), Soilborne flag smut (<i>Urocystis agropyri</i>)	All States	100 mL	This product may be applied undiluted or diluted with up to 500 mL of water per 100 kg of seed. Seed must be treated to ensure even coverage of all grains, using correctly calibrated seed treatment equipment. Reduced emergence may occur if seed treated with flutriafol is sown to a depth greater than 5 cm or into soil at a temperature less than 5°C or greater than 20°C.
Barley	Covered smut (<i>Ustilago segetum</i> var <i>segetum</i>), Loose smut (<i>Ustilago segetum</i> var <i>tritici</i>)			
Oats	Covered smut (<i>Ustilago segetum</i> var <i>segetum</i>), Loose smut (<i>Ustilago segetum</i> var <i>avenae</i>)			
Triticale	Loose smut (<i>Ustilago segetum</i> var <i>tritici</i>)			
Wheat, Barley, Oats and Triticale	Protection against insect pests of stored grains: Lesser grain borer (<i>Rhyzopertha dominica</i>), Granary weevil (<i>Sitophilus granarius</i>), Rice weevil (<i>Sitophilus oryzae</i>), Rust-red flour beetle (<i>Tribolium castaneum</i>), Saw toothed grain beetle (<i>Oryzaephilus surinamensis</i>), Tropical warehouse moth (<i>Ephestia cautella</i>)			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: DO NOT GRAZE OR FEED LIVESTOCK ON TREATED CROPS FOR 4 WEEKS AFTER PLANTING

GENERAL INSTRUCTIONS

Seed may be treated with Vincit C Seed Dressing Flowable at any time from harvesting to sowing. Treated seed may be held over for sowing in the following season. The product contains cypermethrin for protection against insect pests of stored grain.

FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	---	-----------

Vincit C Seed Dressing Flowable is a member of the DMI group of fungicides. For fungicide resistance management Vincit C Seed Dressing Flowable is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Impact Endure and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Impact Endure or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Cheminova Australia Pty Ltd accepts no liability for any losses that may result from the failure of Impact Endure to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

FUNGICIDE RESISTANCE WARNING

GROUP	C	FUNGICIDE
-------	---	-----------

Vincit C Seed Dressing Flowable is a member of the DMI group of fungicides. For fungicide resistance management Vincit C Seed Dressing Flowable is a Group C fungicide. Some naturally occurring individual fungi resistant to Vincit C Seed Dressing Flowable and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Vincit C Seed Dressing Flowable and other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Cheminova Australia Pty Ltd accepts no liability for any losses that may result from the failure of Vincit C Seed Dressing Flowable to control resistant fungi.

MIXING

Vincit C Seed Dressing Flowable may be applied neat at 100mL/100kg of seed or diluted with up to 500 mL of water/100 kg of seed (total application volume 600 mL/100 kg). Calculate the amount of water required (see mixing table). Add half the volume to the tank, then the required amount of Vincit C Seed Dressing Flowable. Make up the final volume with water and mix thoroughly. If allowed to stand for long periods eg. overnight, agitate thoroughly before using. DO NOT allow to stand for more than 24 hours.

APPLICATION VOLUME PER 100 kg OF SEED	VOLUME OF WATER	VOLUME OF VINCIT
100 mL	-	100 mL
200 mL	100 mL	100 mL
300 mL	200 mL	100 mL
400 mL	300 mL	100 mL
500 mL	400 mL	100 mL
600 mL	500 mL	100 mL

APPLICATION

Choose the lowest application volume necessary to give full coverage of seed with the equipment to be used. Application volumes greater than 600 mL are likely to result in excessive wetting of the seed. To achieve disease control, it is essential that the correct quantity of Vincit C Seed Dressing Flowable is applied to the grain. It is also essential that each grain is properly treated.

Undiluted

It is essential to use accurately calibrated equipment which is designed for proper coverage of seed at low application volumes.

Diluted

Use correctly calibrated seed treatment equipment to achieve proper grain coverage.

COMPATIBILITY

Vincit C Seed Dressing Flowable should be applied to grain alone and not mixed with any other product.

CROP SAFETY

Do not apply 'Vydate' to seed pre-treated with Vincit C Seed Dressing Flowable.

CLEANING UP

Equipment used should be cleaned with water. A detergent may be used to assist in the removal of dry deposits.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Short coleoptile cultivars should not be sown too deep. In some situations trifluralin herbicides and triazole fungicides interact to suppress emergence, if possible avoid this contamination. Any seed not intended to be used for sowing should be destroyed.

PROTECTION OF LIVESTOCK

Seed treated with this product must not be used for animal consumption or be allowed to contaminate seed intended for animal consumption. Seed should not be left exposed on the ground (particularly during sowing). Destroy treated seed not intended for sowing.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed or expose treated seed to wildlife and domestic animals, particularly birds. Harmful to fish. DO NOT contaminate streams, rivers or waterways with the chemical or containers which have held the chemical or treated seed.

STORAGE AND DISPOSAL (10L, 200L)

Seed treated with this product must not be used for human consumption or be allowed to contaminate seed intended for human consumption. Treated seed when stored should be kept apart from other seed and the containers should be clearly marked to indicate the contents have been treated with this product. Containers which have held treated seed should not be used for any other purpose.

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways desirable vegetation and tree roots. Empty containers and product should not be burnt.

STORAGE AND DISPOSAL (100 and 110L only)

Seed treated with this product must not be used for human consumption or be allowed to contaminate seed intended for human consumption. Treated seed when stored should be kept apart from other seed and the containers should be clearly marked to indicate the contents have been treated with this product. Containers which have held treated seed should not be used for any other purpose.

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When preparing spray and using the product wear elbow-length PVC gloves and face shield. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet available from www.cheminova.com.au.

WARNING

This product must be used, handled and stored strictly as directed in accordance with this label, packaging and other reference material ("Directions"). To the extent permitted by law Cheminova Pty Ltd and its related companies will have no liability for any injury, loss or damage arising from a failure to follow the Directions.

APVMA Approval No: 45861/0306

Batch no:

Date of Manufacture:

140414-T

Bar Code

Drummaster logo (where appropriate for pack size)

® Registered trademark of Cheminova

<p>FOR 24 HOUR SPECIALIST ADVICE IN EMERGENCY ONLY PHONE 1800 033 111</p>
--