

POISON**KEEP OUT OF REACH OF CHILDREN****READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Vincit® C

Seed Dressing Liquid

ACTIVE CONSTITUENTS: 25 g/L FLUTRIAFOL**4 g/L CYPERMETHRIN****SOLVENT: 495 g/L ETHYLENE GLYCOL****GROUP 3 FUNGICIDE****GROUP 3A INSECTICIDE**

Seed dressing treatment for wheat, barley, oats and triticale to protect against smuts and bunt

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT**Contents 1000L**

Cheminova Australia Pty Ltd ABN 59 472 616 773 / ACN 110 199 169

12 Julius Avenue, North Ryde, NSW 2113

Telephone: 1800 624 597 • www.cheminova.com.au

140430-1000F

Batch No:

Date of Manufacture

**CHEMINOVA**
HELPING YOU GROW

DIRECTIONS FOR USE

RESTRAINT: Do NOT apply 'Vydate' to seed pretreated with this product.

Crop	Disease Controlled	State	Rate/ 100 kg seed	Critical Comments
Wheat	Bunt (<i>Tilletia foetida</i> , <i>T. caries</i>), Loose smut (<i>Ustilago segetum</i> var <i>tritici</i>), Soilborne flag smut (<i>Urocystis agropyri</i>), Seedborne flag smut (<i>Urocystis agropyri</i>)	All States	100 mL	This product is to be applied to grain undiluted. Seed must be treated to ensure even coverage of all grains, using correctly calibrated seed treatment equipment. Reduced emergence may occur if seed treated with flutriafol is sown to a depth greater than 5 cm or into soil at a temperature less than 5°C or greater than 20°C.
Barley	Covered smut (<i>Ustilago segetum</i> var <i>segetum</i>), Loose smut (<i>Ustilago segetum</i> var <i>tritici</i>)			
Oats	Covered smut (<i>Ustilago segetum</i> var <i>segetum</i>), Loose smut (<i>Ustilago segetum</i> var <i>avenae</i>)			
Triticale	Loose smut (<i>Ustilago segetum</i> var <i>tritici</i>)			
Wheat, Barley, Oats and Triticale	Protection against insect pests of stored grains: Lesser grain borer (<i>Rhyzopertha dominica</i>), Granary weevil (<i>Sitophilus granarius</i>), Rice weevil (<i>Sitophilus oryzae</i>), Rust-red flour beetle (<i>Tribolium castaneum</i>), Saw toothed grain beetle (<i>Oryzaephilus surinamensis</i>), Tropical warehouse moth (<i>Ephestia cautella</i>)			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

DO NOT GRAZE PLANTS GROWN FROM TREATED SEED, OR CUT FOR STOCK FOOD, WITHIN 4 WEEKS OF SOWING

GENERAL INSTRUCTIONS

Seed may be treated with Vincit C Seed Dressing Liquid at any time from harvesting to sowing. Treated seed may be held over for sowing in the following season. This product contains cypermethrin for protection against insect pests of stored grain. THIS PRODUCT IS NOT MISCIBLE WITH WATER AND CANNOT BE DILUTED. Equipment must be dry.

FUNGICIDE RESISTANCE WARNING

GROUP 3 FUNGICIDE

Vincit C Seed Dressing Liquid is a member of the DMI group of fungicides. For fungicide resistance management Vincit C Seed Dressing Liquid is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Impact Endure and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Impact Endure or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Cheminova Australia Pty Ltd accepts no liability for any losses that may result from the failure of Impact Endure to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

INSECTICIDE RESISTANCE WARNING

GROUP 3A INSECTICIDE

For insecticide resistance management Vincit C Seed Dressing Liquid is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Vincit C Seed Dressing Liquid and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Vincit C Seed Dressing Liquid or other Group 3A insecticides are used repeatedly. The effectiveness of Vincit C Seed Dressing Liquid on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Cheminova Australia Pty Ltd. accepts no liability for any losses that may result from the failure of Vincit C Seed Dressing Liquid to control resistant insects.

Cheminova Bifenthrin 250EC may be subject to specific resistance management strategies. For further information contact your local supplier, Cheminova representative or local agricultural department agronomist.

CLEANING UP

Equipment should be cleaned with methylated spirits, not water.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Short coleoptile cultivars should not be sown too deep. In some situations trifluralin herbicides and triazole fungicides interact to suppress emergence, if possible avoid this contamination. Any seed not intended to be used for sowing should be destroyed.

PROTECTION OF LIVESTOCK

Seed treated with this product must not be used for animal consumption or be allowed to contaminate seed intended for animal consumption. Seed should not be left exposed on the ground (particularly during sowing). Destroy treated seed not intended for sowing.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed or expose treated seed to wildlife and domestic animals, particularly birds. Harmful to fish. DO NOT contaminate streams, rivers or waterways with the chemical, or containers which have held the chemical or treated seed.

STORAGE AND DISPOSAL

Seed treated with this product must not be used for human consumption or be allowed to contaminate seed intended for human consumption. Treated seed when stored should be kept apart from other seed and the containers should be clearly marked to indicate the contents have been treated with this product. Containers which have held treated seed should not be used for any other purpose. Store product in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.

Non-refillable containers only

Triple or preferably pressure rinse empty containers before disposal. Dispose of rinsings in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. If recycling, replace cap and return containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

100L, 110L

DO NOT remove or tamper with the dry valves or security seal. DO NOT contaminate the drum with water or any other foreign matter. After each use of the product ensure that the dry valve coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the drum is empty remove the dry valve coupler and return to the point of purchase.

Other refillable containers (eg 1000L)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if swallowed. May irritate the eyes and skin. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Do not inhale vapour, dust or spray mist. When preparing and using the product wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet available from www.cheminova.com.au.

WARNING

This product must be used, handled and stored strictly as directed in accordance with this label, packaging and other reference material ("Directions"). To the extent permitted by law Cheminova Pty Ltd and its related companies will have no liability for any injury, loss or damage arising from a failure to follow the Directions.

APVMA Approval No: 30255/0306

Batch no:

Date of Manufacture:

140414-T

Bar Code

Drummaster logo (where appropriate for pack size)

® Registered trademark of Cheminova

**FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111**