

Put Canopy on early and yield late season rewards



Growers who use Canopy early season are less likely to need a late season admiral. Canopy is not just an adjuvant - it's a registered insecticide in it's own right.

DECEMBER

Use 1L of Canopy each time you spray to suppress aphids and silverleaf whitefly. Canopy will not flare mites or whitefly.

JANUARY-FEBRUARY

If mirids appear, add 1L Canopy to low label rates of Shield[®], Transform[®] or Steward[®]. You'll also suppress silverleaf whitefly if they're present. A need for expensive late season sprays is unlikely.*

MARCH-APRIL

Apply Canopy at 0.5L with your defoliant of choice.

10 Reasons to have Canopy in your shed from the start of the season

- Registered for silverleaf whitefly suppression
- Registered for green mirid control
- Registered for aphid control
- Does not flare mites or whitefly
- An adjuvant for other insecticides
- Convenient and simple to apply in every tank mix
- Low rate high performing defoliant aid
- Low rate herbicide adjuvant
- Has no IRMS restrictions
- Registered insecticide uses in canola, chickpeas and pulses.



*excluding ex-field influxes

© Canopy is a registered trademark of Caltex Australia Petroleum Pty Ltd.

This publication is a guide only and no substitute for professional advice. Always read the label before use. FMC Crop Protection Pty Ltd bears no responsibility for the information contained within this publication. Product labels are available at fmc.crop.com.au
TN-INS-16-03