

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FMC

IMAZAPIC

240 HERBICIDE



ACTIVE CONSTITUENT: 240 g/L IMAZAPIC
present as the AMMONIUM SALT

GROUP B HERBICIDE

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaved weed in peanuts and sugarcane as specified in the DIRECTIONS FOR USE table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

FMC

FMC Crop Protection Pty Ltd
5 Palmer Place
Murarrie QLD 4172
Phone: 1800 066 355

Contents **20L**

DIRECTIONS FOR USE**RESTRAINTS:**

DO NOT apply to soils saturated with water.

DO NOT apply by aircraft.

Situation	Weeds Controlled	Rate/ha	Critical Comments
<p>Fallow prior to planting winter crops as directed in the follow crops table (Qld and Northern NSW only)</p>	<p>Barnyard grass (<i>Echinochloa colona</i>), Liverseed grass (<i>Urochloa panicoides</i>), Stink grass (<i>Eragrostis cilianensis</i>), Blowaway grass (<i>Panicum decompositum</i>), Button grass (<i>Dactyloctenium radulans</i>), Pigweed (<i>Portulaca oleracea</i>), Yellow vine (<i>Tribulus terrestris</i>), Mintweed (<i>Salvia reflexa</i>), Boggabri weed (<i>Amaranthus metchelli</i>), Dwarf amaranth (<i>Amaranthus macrocarpus</i>), Peachvine/cowvine (<i>Ipomoea lonchophylla</i>)</p>	<p>150 to 200mL</p>	<p>Treatments may be applied from the end of the previous crop harvest but not less than 3 months prior to planting of winter crop (refer to follow crops table). Best results are obtained where treatments are applied to dry, weed-free soil prior to weed germination. Treatments can be applied to standing crop residues. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5 cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Emerging weeds must be controlled by prior cultivation, or by application of a knockdown herbicide. Where infestation of emerged weeds is light, tank mix treatment with glyphosate or paraquat. Cultivation following application of treatments may reduce pre-emergence weed control provided by FMC Imazapic 240 Herbicide. Prolonged wet soil conditions following treatment will reduce the effective life of pre-emergence treatments. Weed escapes may require follow up application of knockdown herbicides. DO NOT apply more than 200 mL/ha per season on cropland. DO NOT apply in more than two successive seasons on cropland. DO NOT overspray headlands.</p>
<p>Sugarcane (Qld, NSW and WA only)</p>	<p>Awnless barnyard grass (<i>Echinochloa colona</i>), Barnyard grass (<i>E. crusgalli</i>), Blackberry nightshade (<i>Solanum nigrum</i>), Blue billygoat weed (<i>Ageratum houstonianum</i>), Common sida (<i>Sida rhombifolia</i>), Giant/Black pigweed (<i>Trianthema portulacastrum</i>), Glossy nightshade (<i>Solanum americanum</i>), Green amaranth (<i>Amaranthus viridis</i>), Green summer grass (<i>Brachiaria subquadriflora</i>), Ipomoea spp., Guinea grass (<i>Panicum maximum</i>), Liverseed grass (<i>Urochloa panicoides</i>), Milkweed (Mexican fire plant) (<i>Euphorbia heterophylla</i>), Pigweed (<i>Portulaca oleracea</i>), Summer grass (<i>Digitaria ciliaris</i>)</p>	<p>300 to 400mL</p>	<p>DO NOT apply more than once a year to the same crop. Pre-emergence application: PLANT CANE: Apply after planting and before emergence of the cane. On sandy soil, ensure thorough coverage of sets to a minimum depth of 5 cm. RATOON: Apply after harvest and before new cane growth appears. Use the higher rates of FMC Imazapic 240 Herbicide for increased residual control of weeds or anticipated high weed pressure. Heavy trash layers may prevent the product from reaching the soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5 cm. Emerging weeds must be controlled by prior cultivation, or application of an appropriate herbicide. Weed control can be improved under dry conditions or where incorporating moisture has been insufficient, by cultivating the treated soil above sets. Care needs to be taken to minimize exposure of untreated soil or sets. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. Heavy rain and/or irrigation within 2 days following application may concentrate FMC Imazapic 240 Herbicide in the furrow and cause temporary yellowing and stunting of cane leaves. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Weed escapes may require follow up application of an appropriate herbicide. Contact FMC Crop Protection Pty Ltd for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label)</p>

Situation	Weeds Controlled	Rate/ha	Critical Comments
Sugarcane (Qld, NSW and WA only)	As per previous page plus Crowsfoot grass (<i>Eleusine indica</i>)	400 mL plus 3L Pendimethalin 330EC OR 2.2L Pendimethalin 455	Incorporate into the soil within 7 days of application, either by overhead irrigation, rainfall or mechanical incorporation, to a minimum depth of 5 cm. Read all directions for use in sugarcane as well as the Restraints and Follow Crops restrictions on the Pendimethalin 330E or Pendimethalin 455 label.
	As above plus control of those weeds expected with these rates of 250 g/L paraquat products alone.	300 to 400 mL plus 1.2 to 1.6L of 250 g/L paraquat	Post-emergence application (in emerged cane): ALWAYS tank mix FMC Imazapic 240 Herbicide with registered and appropriate rates of paraquat for the emerged weeds. Use the higher rates of FMC Imazapic 240 Herbicide for increased residual control of weeds or anticipated high weed pressure. Use the higher rates of paraquat for dense, more mature weeds. Apply as a carefully directed inter-row spray to weeds and at the base of the plant, minimizing contact with sugarcane foliage. Contact with sugarcane foliage may cause temporary crop damage observed as inter-veinal yellowing and reduction in crop biomass or crop height. Refer to Paraquat (or equivalent) label re spray book types and precautions. Reduce the incidence of "fines" by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis. DO NOT cultivate within 1 hour of application. Heavy rain and/or irrigation within 2 days following application may concentrate FMC Imazapic 240 Herbicide in the furrow and cause temporary yellowing and stunting of cane leaves. DO NOT spray plants covered with heavy dew. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. Contact FMC Crop Protection Pty Ltd for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label).

Situation	Weeds Controlled	Rate/ha	Critical Comments
Peanuts (Qld, NSW and NT only)	Awnless barnyard grass (<i>Echinochloa colona</i>), Barnyard grass (<i>E. crus-galli</i>), Blackberry nightshade (<i>Solanum nigrum</i>), Blue billygoat weed (<i>Ageratum houstonianum</i>), Common sida (<i>Sida rhombifolia</i>), Giant/Black pigweed (<i>Triathema portulacastrum</i>), Glossy nightshade (<i>Solanum americanum</i>), Green amaranth (<i>Amaranthus viridus</i>), Green summer grass (<i>Brachiaria subquadriflora</i>), Ipomoea spp., Guinea grass (<i>Panicum maximum</i>), Liverseed grass (<i>Urochloa panicoides</i>), Milkweed (Mexican fire plant) (<i>Euphorbia heterophylla</i>), Pigweed (<i>Portulaca oleracea</i>), Summer grass (<i>Digitaria ciliaris</i>)	300 to 400 mL	Pre-emergence application. Optimum weed control will occur when moisture is maintained in the top 5 cm of soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5 cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. DO NOT apply to any one crop more than once per season.
	As above plus Nutgrass* (<i>Cyperus rotundus</i>)	400 mL plus Hasten ^{TM1} or Kwicken ^{TM2} at 1L/100L (or a similar esterified oil adjuvant)	Post-emergence application. Apply prior to crop flowering. DO NOT apply to any one crop more than once per season. *Nutgrass: suppression only when applied to nutgrass at the 3 to 4 leaf stage.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING/STOCKFEED WHP:

WHEAT, BARLEY, CHICKPEA, PEANUTS: DO NOT GRAZE TREATED VEGETATION, STUBBLE OR WEEDS, OR CUT FOR STOCKFOOD, FOR 4 WEEKS AFTER APPLICATION.

SUGARCANE: DO NOT GRAZE CROP OR SPRAYED VEGETATION, OR CUT FOR STOCKFOOD, FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

All Applications

Reduce the incidence of “fines” by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis.

Fallow application

FMC Imazapic 240 Herbicide can be applied in fallow prior to the planting of various crops as outlined in the Follow Crops table. FMC Imazapic 240 Herbicide is a pre-emergence herbicide, which is recommended to provide short term (6 to 12 weeks) pre-emergence control of summer germinating grasses and broadleaved weeds in cultivated or uncultivated fallow. A green weed cover of greater than 50% will significantly reduce the length of pre-emergence weed control.

Sugarcane application

Pre-emergence

Where possible, apply to soil two days prior to incorporation. Incorporate with adequate irrigation or rainfall to a minimum depth of 5 cm. Incorporation by flood irrigation may result in loss of weed control. FMC Imazapic 240 Herbicide should be applied to plant or ratoon cane before the cane has emerged. When applying FMC Imazapic 240 Herbicide in moist conditions, allow weeds to germinate prior to application and apply mixed with an appropriate knockdown herbicide. Whenever emerged weeds are present, FMC Imazapic 240 Herbicide should be mixed with an appropriate knockdown herbicide.

Post-emergence

FMC Imazapic 240 Herbicide MUST always be mixed with registered rates of paraquat and applied as a carefully directed inter-row spray to weeds and at the base of the sugarcane stalks, minimizing spray contact with sugarcane foliage. DO NOT add crop oil to this mixture. Other surfactant or adjuvant is not required.

Peanut application

Pre-emergence:

Apply to soil and incorporate with adequate irrigation or rainfall to a minimum depth of 5 cm.

Post-emergence:

FMC Imazapic 240 Herbicide must always be mixed with an adjuvant as directed in the DIRECTIONS FOR USE section of the label.

Soil Type:

Pre-emergence weed control with FMC Imazapic 240 Herbicide can be limited when applied onto soils with pH <5.0 and/or which contain high concentrations of iron (Fe) and/or aluminium (Al). These situations are common in soil types such as Krasnozems and red brown Ferrosols.

MIXING

FMC Imazapic 240 Herbicide is an aqueous solution. Part fill the spray tank with water, then with agitator running, add the required amount of product, then fill the tank with water. When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product. For post-emergence applications in peanuts, add adjuvant after the addition of FMC Imazapic 240 Herbicide to the tank.

No adjuvants or surfactants are required when FMC Imazapic 240 Herbicide is used alone for pre-emergence weed control or with paraquat. DO NOT add crop oil when applied post-emergence with paraquat.

APPLICATION

DO NOT apply by aircraft.

Fallow application:

Apply with boom equipment in not less than 50 L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands.

Sugarcane:

Ensure thorough coverage of soil and weeds with well-calibrated spray equipment. Apply in a minimum of 200 L water/ha using a nozzle configuration to generate a medium to coarse spray pattern for pre-emergence applications or medium spray pattern for post-emergence applications. Replace worn or old nozzles and calibrate equipment on a regular basis.

Peanuts:

Apply with boom equipment in not less than 50 L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands. For post-emergence applications, FMC Imazapic 240 Herbicide should be applied a minimum of 2 hours before rainfall.

EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following the use of FMC Imazapic 240 Herbicide and before use with other products.

COMPATIBILITY

FMC Imazapic 240 Herbicide is compatible with Pendimethalin 330, Pendimethalin 455, atrazine, glyphosate, paraquat, paraquat + diquat, Metsulfuron, Triclopyr 600, Touchdown+ and 2,4-D.

It is NOT compatible with dicamba.

GROUP	B	HERBICIDE
--------------	----------	------------------

RESISTANT WEEDS WARNING

FMC Imazapic 240 Herbicide is a member of the imidazolinone group of herbicides. The product has the inhibitors-of-acetolactate-synthase (ALS) mode of action. For weed resistance management, the product is a Group B herbicide. Some naturally-occurring weed biotypes resistant to this product and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population, if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group B herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, FMC Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

FOLLOW CROPS

Under conditions which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible. The following minimum re-cropping intervals (months after application) should be observed.

Following use in summer fallow:-

MONTHS AFTER APPLICATION				
3	4	10	24	36
Faba beans, Peanuts, mungbeans, maize varieties with CLEARFIELD technology ONLY : -Pacific Hycorn 62IT -Pacific Hycorn 53IT -Pioneer 3395IR; wheat varieties with CLEARFIELD Technology only.	Wheat* (except for wheat varieties with CLEARFIELD Technology ONLY), barley*, lucerne*, chickpea	Sorghum*, maize* (except for maize varieties with CLEARFIELD Technology ONLY), dryland cotton*	Irrigated cotton, sunflower	Any other crops

* The following additional requirements apply if it is intended to plant wheat (except for varieties with CLEARFIELD Technology), barley or lucerne during the next winter season.

DO NOT apply FMC Imazapic 240 Herbicide in fallow later than the end of December.

DO NOT apply FMC Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 200 mm.

= The following additional requirements apply if it is intended to plant sorghum or maize (except for varieties with CLEARFIELD Technology) during the next summer season.

DO NOT apply FMC Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of sorghum or maize is expected to be below 500 mm.

+ The following additional requirements apply if it is intended to plant dryland cotton during the next summer season.

DO NOT apply FMC Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of dryland cotton is expected to be below 550 mm.

Following use in sugarcane and peanuts:-

MONTHS AFTER APPLICATION						
0	4	5	10	18	24	36
Mung beans, Peanuts, soybeans, maize varieties with CLEARFIELD Technology ONLY : -Pacific Hycorn 62IT -Pacific Hycorn 53IT -Pioneer 3395IR	Chick peas, lucerne*, barley*, wheat* (except for wheat varieties with CLEARFIELD Technology ONLY)	Lupins, pasture legumes	Maize* (except varieties with CLEARFIELD Technology ONLY), dryland cotton*	Sorghum*	Irrigated cotton, sunflower	Any other crops

*The following additional requirements apply if it is intended to plant wheat (except for varieties with CLEARFIELD Technology), barley or lucerne during the next winter season.

DO NOT apply FMC Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 200 mm.

=The following additional requirements apply if it is intended to plant sorghum or maize (except for varieties with CLEARFIELD Technology), during the next summer season.

DO NOT apply FMC Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of sorghum or maize is expected to be below 500 mm.

+The following additional requirements apply if it is intended to plant dryland cotton during the next summer season.

DO NOT apply FMC Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of dryland cotton is expected to be below 550 mm.

RE-ENTRY PERIOD FOR FMC IMAZAPIC 240 HERBICIDE PLUS Pendimethalin 455

When tank-mixing FMC Imazapic 240 Herbicide with Pendimethalin 455, DO NOT allow entry into treated areas for 12 hours. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT spray within 50 m of wetlands or waterways.

CROP SAFETY

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth in sugarcane but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under conditions where soil moisture is high within 2 days following application and on light/sandy soil. Contact FMC Australasia Pty Ltd for advice on variety susceptibility before using this product.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Keep out of reach of children.

Store in the closed, original container, in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or preferably pressure rinse containers before disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection point or similar container management program site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves.

When tank-mixing with Pendimethalin 330EC, Pendimethalin 455 or Paraquat 250, consult also the safety directions for those products.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information on Imazapic 240 WG Herbicide refer to the Material Safety Data Sheet (MSDS) which is available from the supplier or from fmccrop.com.au.

WARRANTY

FMC Crop Protection Pty Ltd makes no warranty expressed or implied, concerning the use of this product other than that indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

APVMA Approval No: 69495/60935

® FMC is a Registered Trademark of FMC Corporation Pty Ltd.



FMC Crop Protection Pty Ltd.
A.B.N. 48 159 288 123
5 Palmer Place
Murarrie QLD 4172
Phone: 1800 066 355
www.fmccrop.com.au
Technical Enquiries: 1800 901 939

In a Transport
Emergency Dial
000
Police or Fire
Brigade

SPECIALIST ADVICE
IN EMERGENCY ONLY
1800 033 111
ALL HOURS –
AUSTRALIA WIDE

