

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FMC 2,4-D LV Ester

680 HERBICIDE

ACTIVE CONSTITUENT: 680 g/L 2,4-D present
as the 2-ethylhexyl ester

GROUP

HERBICIDE

A low volatile herbicide for selective control of various weeds
in crops, pastures and non-agricultural areas as
specified in the DIRECTIONS FOR USE table.

This is a PHENOXY HERBICIDE that can cause severe damage
to susceptible crops. Please read the Drift Warning Statement.

IMPORTANT: READ THIS LEAFLET BEFORE USE

FMC

FMC Crop Protection Pty Ltd
Unit 28/3 Metroplex Avenue,
Murarrie Qld 4172
Phone: 1800 901 939

DIRECTIONS FOR USE**RESTRAINTS**

AVOID spraying if rain is likely within 6 hours or if strong winds prevail.

DO NOT apply if crop of weeds are stressed due to dry or excessively moist conditions.

1. FIELD CROPS

Situation & Crop	Weeds Controlled	State	Rate /ha	Critical Comments
Wheat, Barley	Refer to Weed Table	Vic only	210 - 800 mL	CROP STAGES: ALL CEREALS Vic only: Apply at tillered to boot stages. NSW only: Apply after when the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller. Qld only: Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling of the head at the top of the main stem). SA, Tas only: Apply from completion of tillering to early jointing stage.
		SA only	230 - 800 mL	
		Qld, NSW, ACT only	405 - 800 mL	
		Tas only	620 - 800 mL	
		WA only	800 mL	
Triticale		Qld, NSW, ACT only	405 - 800 mL	
		SA only	240 - 820 mL	
		Vic only	210 - 800 mL	
Cereal Rye		Qld, NSW, ACT only	405 - 800 mL	
		Vic only	210 - 800 mL	
Sugar Cane		Qld only	1.15 - 2.4 L	Post-emergence.
Stubble/Fallow Spray Prior to Direct Drilling or Sowing. Winter Cereals, Grain Legumes (Peanuts Qld only), Canola		ALL STATES	210 - 800 mL	Observe the plant back periods given in the table in this leaflet. Must be tank mixed with a knockdown herbicide such as FMC Glyphosate 450 Herbicide, paraquat or paraquat/diquat mixtures. Select appropriate rate from the weed table. For Skeleton Weed, spraying should only be done 6 - 8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.

Situation & Crop	Weeds Controlled	State	Rate /ha	Critical Comments
Harvest Aid or Salvage Spray - Winter Cereals	Broadleaf Weeds Refer to Weed Table	ALL STATES	1.68 L	Apply after dough stage of crop. Interval between application and effectiveness is 10 - 20 days. For desiccation of green matter, estimate harvest date and apply spray approximately 14 days earlier. Rain between spraying and actual harvest can negate results. Note: Where thistles are tall and branching above the crop, spraying can turn the branches down into the crop, presenting more stalks to cause header comb blockages. Spraying may increase seed contamination of harvest by accelerating maturity. DO NOT use with undersown legumes that have not set seed.
Potatoes Pre-harvest Preparation	Broadleaf Weeds such as Clover, Variegated Thistle and Cruciferous weeds	Vic, Tas only	1.15 - 2.4 L	Apply approximately 4 to 5 weeks before harvest after the potato haulms have dried off. Use the highest rate where weeds are more than 30 cm in height. For boom spraying at least 100 litres of spray mixture per hectare. If grasses such as ryegrass and Winter grass are also present add amitrole.

2. PASTURE, NON-AGRICULTURAL, INDUSTRIAL

Situation & Crop	Weeds Controlled	State	Rate /ha	WHP (days)	Critical Comments
Improve Pasture containing Clovers	Refer to Weed Table	Qld, NSW, ACT, Tas, SA only	405 – 800 mL	7	Clover must be well covered by the grass or extensive damage may result.
Pastures – Non legumes, Rights of Way & Industrial		Qld, NSW, ACT, Tas, SA only	800 mL - 4.68 L		Control of most perennial weeds, but due to the rooting habits of most species control may take a number of years. Damage may result to legumes in pasture.
Pastures – Direct Drilling or Surface Sowing	Charlock, Clover, Medics, Mustards, Paterson's Curse, Saffron, Slender, Variegated and Spear Thistles, Turnip Weed, Wild Radish, Wild Turnip	NSW only	800 mL - 1.5 L (Aerial Application)	Applying to young, actively growing weeds. Sowing: DO NOT sow pasture seed for at least 21 days after application. If soil moisture is dry, delay sowing for at least 30 days.	
			800 mL - 6.5 L		
		70 – 620 mL/100 L			
		800 mL - 1.5 L (Ground Application)			
	As Above plus: Capeweed, Wireweed, Storksbill/ Erodium, Flatweed, Horehound (seedlings), Skeleton Weed, Nodding or Star Thistles.		800 mL - 1.5 L (Ground Application)		
	St John's Wort		3.24 - 4.68L (Aerial or Ground)		
	All of above plus grasses		As above plus glyphosate		

3.SPOT SPRAYING

Situation & Crop	Weeds Controlled	State	Rate/ha	WHP (days)	Critical Comments
Spot Spraying (All Situations)	Refer to Weed Table	All States	1/100 th of rate on Weed Table per 10 L water per 100 m ²	7	Apply through Knapsack. Thorough wetting of weed is essential.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15 APRIL TO 15 SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WITHHOLDING PERIOD (WHP):

PASTURES, CEREAL CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED

WEED TABLE

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the crop in the table below. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed e.g. *Rumex* spp. (Docks) and *Polygonum* spp. (Wireweed, Climbing Buckwheat) are killed to ground level only.

APPLICATION RATE PER HECTARE

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
Amaranthus spp.	-	-	-	800 mL	-	-	-	-	
Angled Onion	-	-	-	-	-	-	3.26 L	0.8 - 1.68 L	Spray when buds are forming or early flowering

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
Apple Sodom	-	-	-	-	-	-	-	2.9 - 3.26 L	
Bathurst Burr	-	-	-	800 mL	-	-	1.7 - 3.26 L	1.7 - 3.26 L	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Black Knapweed	-	-	-	-	-	-	3.26 L	-	Spray before flowering. DO NOT cultivate these infestations.
Buffalo Burr	-	-	-	-	-	-	-	800 mL - 1.5 L (Not Qld & WA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
California Burr	-	-	-	800 mL	-	-	1.7 - 3.26 L	1.15 - 1.68 L (Not SA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Caltrop	-	-	-	620 mL – 800 mL	-	-	1.7 - 3.26 L	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Cape Tulip	-	-	-	-	-	1.15 L	3.26 L	1.7 - 3.26 L	Spray before flowering.
Capeweed	800 mL	800 mL	800 mL	530 – 800 mL	-	-	-	2.47 - 3.26 L	Spray up to rosette stage.
Charlock	405 mL	405 mL	800 mL	405 mL	-	-	-	800 mL	Spray up to rosette stage.

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
Clover	-	-	-	620 – 800 mL	-	-	-	800 mL	
Colocynth	-	-	-	-	-	-	3.26 L	-	Spray at seedling stage only.
Deadnettle	-	-	-	800 mL	-	-	-	-	
Devil's Claw	-	-	-	800 mL	-	-	1.32 L	-	Spray prior to pods forming.
Dock	800 mL	800 mL	-	-	800 mL	800 mL	-	1.68 - 2.47 L	Spray at rosette stage to kill top growth only.
Fat Hen	-	-	-	405 - 800 mL	-	-	-	-	
Flatweed	-	-	-	800 mL	-	-	-	2.47 - 3.26 L	
Fumitory (red)	-	800 mL	-	800 mL	-	-	-	2.47 - 3.26 L	Spray up to rosette stage.
Fumitory (white)	800 mL	405 mL	-	800 mL	-	-	-	2.47 - 3.26 L	Spray up to rosette stage.
Galvanised Burr	-	-	-	-	-	-	4.68 L	4.68 L (Not Qld & WA)	Spray from seedling to pre-flowering.
Goosefoots	-	-	-	800 mL	-	-	-	-	

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
Hard Head or Russian Knapweed	-	-	-	-	-	-	3.26-5.2 L	-	Spray before flowering.
Hogweed, Wireweed	800 mL	800 mL	-	800 mL	800 mL	-	-	1.15 - 1.68 L (Not SA)	Spray up to rosette stage.
Hoary Cress, Whiteweed	-	800 mL	800 mL	800 mL	-	-	1.7 - 3.26 L	1.7 - 2.1 L	Spray from late rosette to pre-flowering.
Horehound (seedlings)	-	800 mL	-	-	-	840 mL	-	1.7 - 3.26 L	Late Autumn to early Spring.
Ironweed, Corn Gromwell	-	-	-	-	800 mL	-	-	1.15 - 1.68 L	
Khaki Weed	-	-	-	-	-	-	-	800 mL - 1.15 L (Not SA)	Spray young seedlings only.
Lincoln Weed	-	800 mL	-	-	-	-	-	-	Autumn spray before sowing improves control.
London Rocket	-	-	-	-	-	570 mL	-	1.6 - 2.5 L (WA only)	
Lupins	800 mL	-	-	405 – 800 mL	-	-	-	-	Spray up to rosette stage.

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
Melilotus/ Hexham Scent	800 mL	800 mL	-	-	800 mL	-	-	1.15 - 1.68 L	Spray up to rosette stage.
Melons – camel, paddy	-	-	-	405 - 800 mL	-	-	-	-	
Mustards	330 mL	230 – 800 mL	800 mL	405 - 900mL	620 mL	620 mL	3.26 L	1.68 - 2.47 L	Spray up to rosette stage.
Mexican Poppy	-	2.3 - 3.5 L	-	800 mL	-	840 mL	-	800 mL - 1.15 L (1.1 - 1.5 L WA only)	Spray rosette stage and before flowering.
Mintweed	-	-	-	800 mL	620 mL	-	-	800 mL - 1.15 L	Spray active seedlings only.
Muskweed	800 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
New Zealand Spinach	-	-	-	800 mL	-	-	-	-	
Noogoora Burr	-	-	-	800 mL	-	-	1.68 - 3.26 L	1.68 - 3.26 L	Spray seeling to pre-flowering.
Nut Grass	-	-	-	-	-	-	3.26 - 5.2 L	-	Spray within 4 weeks of foliage emergence, repeat spray necessary.

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
Paterson's Curse	-	-	-	800 mL	-	840 mL	1.7 - 3.26 L	800 mL - 1.68 L (1.15 - 1.5 L WA only)	Spray seedling to rosette stage.
Poppy Wild	405 mL	-	-	-	-	-	-	2.1 - 2.9 L	Spray up to rosette stage.
Ragwort	-	-	-	-	-	-	3.26 L	3.26 L	Spray at rosette to cabbage stage.
Rapeseed	800 mL	-	-	405 - 800 mL	-	-	-	-	Spray up to rosette stage.
Rapistrum	-	-	-	-	-	570 mL	-	840 mL (WA only)	
Rough Poppy	-	405 mL	-	405 - 800 mL	-	-	-	800 mL	Spray young seedlings only.
St. John's Wort	-	-	-	-	-	-	3.26- 5.9 L	3.26- 4.68 L	Spray before flowering. Spray before plants are 40 cm high.
Safflower	-	-	-	405 - 800 mL	-	-	-	-	

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
Sand Mustard/ Sand Rocket	-	-	-	-	-	-	3.26 L	-	Spray before flowering.
Shepherd's Purse	-	-	-	800 mL	-	-	-	-	
Silver Leaf Nightshade	-	-	-	-	-	-	3.26 L	-	Spray at flowering. Fallow land: controls top growth only.
Skeleton Weed	-	800 mL	-	800 mL	-	-	3.26 L	1.15 - 1.68 L	Spray rosettes before aerial growth commences.
Stingless Nettle (Deadnettle)	-	800 mL	800 mL	-	-	-	-	2.12 - 2.47 L	
Stinging Nettle	800 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Stinkwort	-	-	-	800 mL	-	-	1.68 - 3.26 L	1.68 - 3.26 L	Spray younger plants. Use higher rate as plants mature.
Storksbill/ Erodium	-	-	-	800 mL	-	-	-	-	
Sunflower Seedlings	800 mL	-	-	405 - 800 mL	620 mL	-	-	-	Spray multiple leaves.
Thistles: - Golden	-	-	-	-	-	-	3.26 L	3.26 L	Spray at rosette stage,

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
- Nodding	-	-	-	-	-	-	3.26 L	1.15 - 1.68 L	Spray at rosette stage to pre-flowering.
- Saffron	620 mL	800 mL	-	405 - 800 mL	800 mL	800 mL	800 mL - 1.68 L	840 mL - 3.26 L (WA only)	Spray up to rosette stage.
- Sheep	-	-	-	-	-	840 mL	-	840 mL - 3.26 L (WA only)	
- Slender, Shore	-	-	800 mL	800 mL	-	-	1.7 - 3.26 L	800 mL - 3.26 L	Spray at rosette stage.
- Soldier	-	-	-	-	-	-	3.26 L	-	Spray at rosette stage
- Spear	800 mL	-	800 mL	-	-	-	800 mL - 2.5 L	1.15 - 2.1 L	Spraying at seedling to rosette stage. Use higher rate as plants mature (pastures).
- Stemless	-	-	-	-	-	-	3.26 L	2.47 - 3.26 L	Spray at rosette stage to flowering.
- St Barnaby's	-	-	-	-	-	-	-	1.15 - 1.68 L	
- Star	-	-	-	800 mL	-	-	1.68 - 3.26 L	1.15 - 1.68 L	Spraying at seedling to rosette stage. Use higher rate as plants mature.

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
- Variegated	-	-	800 mL	405 - 800 mL	620 mL	-	800 mL - 2.5 L	800 mL - 3.26 L	Spray at rosette stage. Can cause stock poisoning.
Thornapple	-	-	-	405 - 800 mL	-	-	3.26 L	800 mL - 1.68 L	Spray at seedling stage.
Tree Hogweed	800 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Turnip Weed	-	405 mL	-	405 - 800 mL	405 mL	620 mL	-	800 mL	Spray seedling only.
Vetches/Tares	800 mL	620 mL	800 mL	-	-	-	-	-	
Wards Weed	-	405 mL	-	-	-	-	-	-	Spray at seedling stage.
Wild Cabbage	800 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Wild Garlic Only	-	-	-	-	-	-	6.62 L	-	Suppresses aerial growth.
Wild Mignonette	-	-	-	-	-	840 mL	3.26 L	-	Spray at rosette stage.
Wild Mustard	-	-	-	-	-	650 mL	-	1.8 - 2.47 L (WA only)	

Weeds Controlled	Crop						Pasture – Non Legume		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, ACT, Tas, SA, Qld, WA only	
Wild Radish	800 mL	800 mL	800 mL	405 – 800 mL	800 mL	570 mL	-	800 mL (840 mL WA only)	Spray up to rosette stage.
Wild Sage	-	-	-	-	-	-	-	2.47 - 3.26 L	
Wild Teasel	-	-	-	-	-	-	1.68 - 3.26 L	-	Spray at rosette stage. Use higher rate as plants mature.
Wild Turnip	210 mL	230 mL	800 mL	405 – 800 mL	-	450 mL	-	800 mL (840 mL WA only)	Spray up to rosette stage.

Plant back days for FMC 2,4-D LV Ester 680 Herbicide

Crop	Rates		
	Up to 510 mL/ha	510 mL – 1.15 L/ha	1.15 - 1.59 L/ha
Balansa Clover	7	7	10
Barley ¹	1	1	3
Chickpeas ²	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14

Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins ⁴	7	14	21
Medics	7	7	10
Narbon Beans	7	7	10
Navy Beans	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola/Rapeseed ²	14	21	28
Rice	7	7	14
Safflower ²	7	14	21
Sorghum ³	3	7	10
Soybean	14	14	21
Sub-Clover	7	7	10
Sunflower ³	7	10	14
Triticale ¹	1	3	7
Vetch	7	7	10
Wheat ¹	1	3	7
White Clover	7	7	10

IMPORTANT: WHEN APPLIED TO DRY SOILS AT LEAST 15 mm (1/2 inch) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

Notes:

- 1 In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for wheat, barley and triticale.
- 2 In Queensland, planting of canola/rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall at least 15 mm.
- 3 In Central Queensland, when using 730 mL/ha or less of FMC 2,4-D LV Ester 680 Herbicide, the Plant Back Period for sorghum and sunflower is 1 day irrespective of rainfall.
- 4 In WA the Plant Back Period for lupins at all rates is 28 days.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

DO NOT spray in high winds.

DO NOT spray crops or weeds outside the stages indicated in the Critical Comments as damage, loss of yield or inadequate weed control may result.

APPLICATION INFORMATION

This product may be used in either high or low volume sprays. Just pour into water and stir.

Boom Spraying: Use 30 - 100 litres water per hectare.

Aerial Spraying: Use 40 - 90 litres water per hectare.

Note: Refer to the Department of Agriculture/Primary Industries in your state for the current restricted spraying areas.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible. Otherwise wash out the unit with hot soapy water followed by several clear water rinses. DO NOT use wooden spray vats, as they cannot be cleaned.

Hoses cannot be cleaned and new hoses should be fitted when the unit is to be used or any other purpose.

COMPATIBILITY

This product can be tank mixed with atrazine, chlorsulfuron, dicamba, glyphosate, metsulfuron, paraquat, paraquat/diquat, fluroxypyr, triasulfuron and triclopyr.

Note:

1. As formulations of other manufacturer's products are beyond the control of FMC Crop Protection all mixtures should be tested on a small scale before mixing in the spray tank.

2. Tank Mixing Instructions

- Fill the tank ¼ full with water and agitate.
- Add wettable powders and water dispersible granules first.
- Agitate until these are uniformly dispersed, while adding water until the tank is 90% full.
- Add suspension concentrates (flowables) then soluble concentrates. Add emulsifiable concentrates last.
- Top up the tank with water and continue agitation until all the ingredients are properly mixed.
- Observe any mixing sequence instructions specifically stated on the tank mix products.

RESISTANT WEEDS WARNING

GROUP	I	HERBICIDE
--------------	----------	------------------

FMC 2,4-D LV Ester 680 Herbicide is a member of the Phenoxys group of herbicides. FMC 2,4-D LV Ester 680 Herbicide has the Disruptors of plant cell growth mode of action. For weed resistance management FMC 2,4-D LV Ester 680 Herbicide is a Group I Herbicide.

Some naturally-occurring weed biotypes resistant to FMC 2,4-D LV Ester 680 Herbicide and other Group I Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FMC 2,4-D LV Ester 680 Herbicide or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, FMC Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of FMC 2,4-D LV Ester 680 Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Legume Tolerance

If clovers are present, care should be taken to ensure that they have reached the 3 - 4 leaf stage before spraying. Rates above 405 mL of this product per hectare will destroy most clovers, whilst lucerne and medics are susceptible at any strength.

Drift Warning

DO NOT apply under metrological conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

In particular, avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, ornamentals, oil seed and legume crops or other susceptible plants and trees (e.g. Kurrajongs, Belahs, Eucalypts).

- DO NOT use unless wind speed is more than 3 kilometres per hour and less than 15 kilometres per hour as measured at the application site.
- DO NOT apply with smaller than coarse to very coarse spray droplets according to the ASAE S572 definition for standard nozzles.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers, or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

DO NOT store near food, feedstuffs, fertiliser or seed.

The method of disposal of the container depends on the container type. Read the 'STORAGE AND DISPOSAL' instructions on the label that is attached to the container.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear PVC or rubber apron and elbow-length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS) and is available from the supplier.

WARRANTY

FMC Crop Protection Pty Ltd makes no warranty expressed or implied, concerning the use of this product other than that indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

APVMA Approval No: 68240/57817

® FMC is a Registered Trademark of FMC Corporation Pty Ltd.



FMC Crop Protection Pty Ltd
A.B.N. 48 159 288 123
Unit 26, 8 Metroplex Avenue
Murarrie Qld 4172
Phone: 1800 066 355
www.fmccrop.com.au
Technical Enquiries: 1800 901 939

In a Transport
Emergency Dial
000
Police or Fire
Brigade

SPECIALIST ADVICE
IN EMERGENCY ONLY
1800 033 111
ALL HOURS –
AUSTRALIA WIDE