

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FMC Triallate

500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L TRI-ALLATE
SOLVENT: 471 g/L LIQUID HYDROCARBON

GROUP

J

HERBICIDE

For the control of Wild oats in Wheat, Triticale, Chickpeas, Barley, Peas, Linseed, Lupins, Canola (Rapeseed), Faba beans and Safflower as specified in the DIRECTIONS FOR USE table.

IMPORTANT: READ THIS LEAFLET BEFORE USE



FMC

FMC Crop Protection Pty Ltd
5 Palmer Place
Murarrie QLD 4172
Phone: 1800 901 939

DIRECTIONS FOR USE IN ALL STATES

Crop	Weeds	Rate L/ha	Critical Comments
Barley, Triticale, Wheat	Wild Oats	1.6L	Apply immediately prior to or up to 3 weeks before sowing to a friable seedbed under conditions, which will allow for the incorporation of the product to a depth of 2 to 4 cm with either a single or double pass of suitable incorporation equipment (see incorporation). A delay of 1 week between application and planting is recommended for Linseed. COMPLETE AND UNIFORM INCORPORATION OF FMC TRIALLATE 500 IS ESSENTIAL FOR WEED CONTROL. See GENERAL INSTRUCTIONS for details at incorporation.
Chickpeas, Faba beans, Lupins, Peas			FMC Triallate 500 Herbicide may be used in crops undersown with legumes. For shallow-sown legume and oil seed crops, complete incorporation by cultivation to a depth of 5cm must be undertaken before planting.
Linseed, Canola, Safflower			FMC Triallate 500 Herbicide may be tank mixed with chlorsulfuron and triasulfuron for control of annual ryegrass, Paradoxa grass and certain broadleaf weeds prior to planting of wheat only. FMC Triallate 500 Herbicide may be tank mixed with trifluralin (480g/L) for control of annual ryegrass, annual Phalaris, Paradoxa grass, enhanced control of wild oats (emerging from top 1cm of seedbed) and wireweed. Do not apply less than 1 week prior to planting of wheat, barley, triticale and linseed (do not use the tank mix in Qld). Incorporation of this tank mix should be within 4-5 hours of application. Refer to the trifluralin label incorporation details. Read and follow label instructions, restraints, plant-back periods and safety directions for tank mix products.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

**WITHHOLDING PERIOD – HARVEST
NOT REQUIRED WHEN USED AS DIRECTED**

GENERAL INSTRUCTIONS

FMC Triallate 500 Herbicide normally provides 6 to 8 weeks control of wild oats when the seedbed is moist during the emergence of wild oats. Under very dry soil surface (top 2cm) conditions, wild oats may not be controlled. Wild oats may emerge between application and complete incorporation by sowing particularly if harrows provide initial incorporation. FMC Triallate 500 Herbicide may not control wild oats, which have germinated at the time of application and incorporation if the top 2 cm of soil is dry. FMC Triallate 500 Herbicide will not control emerged wild oats.

CROP SAFETY

Reduced emergence of cereals and linseed may occur if seed is placed in treated soil layer. Do not plant cereal crops at greater than 5 to 7 cm. Over spraying of headlands or corners may result in crop damage. On heavy clay soils, crop thinning may occur in wheel tracks. Environmental conditions which adversely affect emergence or early crop growth, may predispose crops to injury, particularly where tank mixes with other pre-emergence herbicides are used.

Read and follow label directions, restraints, plantback periods and safety directions for tank mix products.

SOIL PERSISTENCE AND FOLLOWING CROPS

Under prolonged and extremely dry conditions FMC Triallate may persist for several months. Damage to field oats and sorghum may occur.

MIXING

FMC Triallate 500 Herbicide is an oily concentrate that mixes readily with water. Add the required amount of FMC Triallate 500 Herbicide to half the required amount of water in the spray tank. Mix thoroughly and add the remaining water.

APPLICATION

Apply to soil in friable condition so that uniform incorporation of the product can be achieved to a depth of 2 to 4 cm. The soil should be free of large clods (>75 mm) and should not have crop or weed residue at a level, which will interfere with the free flow of soil through planting equipment providing full soil disturbance. Emerged wild oats should be controlled either by prior cultivation/ application of a knockdown herbicide or by the tank mixing of a knockdown herbicide with FMC Triallate 500 Herbicide.

Apply FMC Triallate 500 Herbicide by ground sprayer in 30 to 100 litres of water per hectare using a boomspray fitted with flat fan nozzles. Use spraying pressure of approximately 200-240kPa. Avoid application under strong and/or gusty wind conditions.

DO NOT apply by aircraft.

INCORPORATION

Two-pass incorporation provides superior weed control to single pass incorporation on heavier textured soils.

- (i) FMC Triallate 500 Herbicide may be incorporated initially in a suitable cultivated seedbed with either heavy 5 row harrows e.g. Phoenix harrow, or tined cultivator. Final incorporation with a combine or airseeder/cultivator is essential to provide for maximum weed control.

Heavy tined harrows or a rotary harrow should trail the planter or follow as a separate operation at an angle to the planter to complete incorporation. Ensure that the soil surface is left free of pronounced ridging.

- (ii) On soils in friable condition, FMC Triallate may be incorporated with the planting operation only up to 24 hours following application onto a dry soil surface or up to 6 hours on moist surface soil. If FMC Triallate has been sprayed on dry soil, and it rains before incorporation is planned, the product must be incorporated within 6 hours to retain its herbicide effectiveness. Planting should be carried out with a combine or airseeder/cultivator that provides full soil disturbance. Heavy tined harrows or a rotary harrow should trail the planter or follow as a separate operation to complete incorporation. Ensure that the soil surface is free of pronounced ridging.

DO NOT operate incorporation equipment at greater than 5 to 7 cm depth.

Where soil is wet at planting, poor incorporation may result. For shallow-sown legume and oil-seed crops, complete incorporation by cultivation to a depth of 5 cm must be undertaken before planting.

Note that point size should be matched to tyne spacing to avoid uneven incorporation. For example, a wideline seeder with 15 cm tyne spacing should be fitted with unworn 15 cm points. A combine with 9 cm spacing should be fitted with an evenly worn set of 10 cm points. Sweeps or other low profile points may not provide satisfactory incorporation.

COMPATIBILITY

FMC Triallate 500 Herbicide is compatible with: chlorsulfuron, triasulfuron, trifluralin and glyphosate.

RESISTANT WEEDS WARNING

FMC Triallate 500 Herbicide is a member of the thiocarbamate group of herbicides. FMC Triallate 500 Herbicide has the inhibition of mitosis mode of action. For weed resistance management FMC Triallate 500 Herbicide is a Group J herbicide.

Some naturally occurring weed biotypes resistant to FMC Triallate 500 Herbicide and Group J herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These weeds will not be controlled by FMC Triallate 500 or other Group J herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use FMC Crop Protection Pty Ltd accepted no liability for any losses that may result from the failure of FMC Triallate 500 Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Dangerous to fish. Do not contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

110L Mini Bulk Returnable Container

Store the original sealed 110L container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Do not tamper with the MicroMatic valve or the security seal. Do not contaminate the container with water or any other foreign matter. After each use of the product ensure the MicroMatic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the 110L container have been used, please return the empty container to the point of purchase. The 110L container remains the property of FMC Crop Protection Pty Ltd.

Refillable Containers: 1000L

Store in the closed, original container in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to point of sale for refill or storage.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed, will irritate eyes and skin, repeated exposure may cause allergic disorders. Do not inhale spray mist, avoid contact with eyes, skin and clothing. When opening the container and using the prepared spray wear cotton overalls, buttoned to the neck and wrist and a washable hat and elbow length PVC gloves, goggles and half facepiece respirator. When using the prepared spray wear cotton overalls buttoned to the neck and wrist. If product on skin, immediately wash area with soap and water, if in eyes wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26. If swallowed DO NOT induce vomiting. Give a glass of water. Avoid giving alcohol.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet that can be obtained from the supplier.

WARRANTY

FMC Crop Protection Pty Ltd makes no warranty expressed or implied, concerning the use of this product other than that indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

APVMA Approval No: 67818/56914

® FMC is a Registered Trademark of FMC Corporation Pty Ltd.



FMC Crop Protection Pty Ltd.
A.B.N. 48 159 288 123
5 Palmer Place
Murarrie QLD 4172
Phone: 1800 066 355
www.fmccrop.com.au
Technical Enquiries: 1800 901 939

In a Transport
Emergency Dial
000
Police or Fire
Brigade

SPECIALIST ADVICE
IN EMERGENCY ONLY
1800 033 111
ALL HOURS –
AUSTRALIA WIDE