



CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

ACTIVE CONSTITUENT:
250 g/L DIFENOCONAZOLE

CONTENTS:
1 – 1000 L

EC FORMULATION TYPE
**Emulsifiable
Concentrate**

SOLVENTS:
580 g/L LIQUID HYDROCARBON
100 g/L N-METHYL-2-PYRROLIDONE

GROUP 3 FUNGICIDE



For the control of target spot of potatoes and tomatoes, leaf blight of carrots, leaf spot diseases of bananas and husk spot on macadamias as specified in the Directions for Use Table.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, **DO NOT** induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet that can be obtained from www.fmccrop.com.au

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

GROUP 3 FUNGICIDE

Bulls-eye® Systemic Fungicide is a member of the DMI Group of fungicides. For fungicide resistance management Bulls-eye® is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Bulls-eye® may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if this fungicide is used repeatedly. These resistant fungi will not be controlled by Bulls-eye®, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FMC accepts no liability for any losses that may result from the failure of Bulls-eye® to control the resistant fungi.

MIXING

Pour the measured volume of Bulls-eye® slowly into the required volume of spray water in the spray tank while maintaining constant agitation. Sprays

containing Bulls-eye® should be used as soon as possible after mixing and maintain constant agitation throughout the spray operation.

WETTING AGENTS

The addition of a non-ionic wetting agent is required when Bulls-eye® is being applied to macadamia nuts and addition of a miscible oil for application to bananas.

COMPATIBILITY

Bulls-eye® is compatible with the following products for the crops in which they are recommended: Sulfos, Rogor, methidathion, Cyren 500EC, Pirimicarb, Permethrin, Alpha-cypermethrin, Ridomil MZ formulations, chlorothalonil, procymidone, iprodione, mancozeb, urea, EC formulations of synthetic pyrethroids, EC formulations of organophosphate insecticides, Copper oxychloride formulations.

SPECIAL WARNING: BANANAS

There are certain conditions when the surface of recently emerged fruit is particularly prone to marking damage from spray applications. In circumstances where application will be made to very rapidly growing fruit in hot conditions with strong direct light. It is recommended that all emerged

bunches be bagged prior to spraying to minimise risk of fruit marking.

If using Bulls-eye® with a mineral oil in blocks with un-bagged bunches, please contact FMC, your chemical reseller or aerial applicator for up to date information regarding the oil formulations recommended by FMC.

APPLICATION

Thorough coverage is essential for effective results.

For aerial application use 20 to 30 L water/ha.

For ground application, use at least 100 L water/ha.

Macadamia Nuts: Apply by high volume (dilute) sprayer or by concentrate sprayer.

Dilute spraying: Use a sprayer designed to apply high volumes of water up to the point of run-off and compatible with the crop being treated. Set up and operate the sprayer to achieve even coverage throughout the crop canopy and apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The sprayer must be calibrated for each individual crop situation to ensure that the correct water volume and crop coverage is achieved.

Concentrate spraying: Use a sprayer designed to apply low volumes of water (less than up to the point of run-off) and compatible with the crop being treated. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your choice of water volume. The water volume used in low volume application is calculated from the amount used in high volume application. The mixture rate for concentrate mixture application can be calculated as follows:

1. If the dilute mixture application spray volume is 1 500 L/ha and your volume choice is 500 L/ha then the concentration factor is $3 \times (1,500\text{L}/500\text{L} = 3)$.
2. If the dilute label rate of application is 50 g/100 L water, then the concentrate rate of application becomes $3 \times 50 \text{ g} = 150 \text{ g}/100 \text{ L}$ of concentrate spray mixture.

The sprayer must be calibrated for each individual crop situation to ensure that the correct water volume and crop coverage is achieved. Users must ensure that they are adequately trained in order to apply Bulls-eye® according to industry guidelines and industry Best Practices.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied to plants at any time.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to aquatic organisms. DO NOT spray within 5 m downwind of water bodies. **DO NOT** apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or waterways. **DO NOT** contaminate

dams, waterways or drains with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

Leaflet

The method of disposal of the container depends on the container type. Read the 'Storage and Disposal' instructions on the label that is attached to the container.

Non-returnable containers only

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

100, 110 L

DO NOT remove or tamper with the dry valves or security seal. **DO NOT** contaminate the drum with water or any other foreign matter. After each use of the product ensure that the dry valve coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the drum is empty remove the dry valve coupler and return to the point of purchase.

1000 L

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

DIRECTIONS FOR USE

RESTRAINT:

DO NOT apply more than 6 applications of this product per season.

The effect of Bulls-eye® could be diminished if rain falls within 2 hours of application.

CROP	DISEASE	STATE	RATE/ha	WHP, Days	CRITICAL COMMENTS
Bananas including Bananas interplanted with Avocados	Yellow Sigatoka (<i>Mycosphaerella musicola</i>) Black Sigatoka (<i>Mycosphaerella fijiensis</i>)	Qld, NSW, NT only	Ground application: 400ml + 3-5 L of water miscible oil* in a convenient volume of water Aerila application: 400 mL + 3-5 L of a water miscible oil* in a minimum of 30 L of water	1	This use is subject to a CropLife Australia Fungicide Resistance Management Strategy. Ground application: Apply by misting machine or airblast sprayer. Use rates towards the higher end of the range where weather conditions favour diseases or where equipment or terrain does not permit thorough spray coverage of all foliage. NSW, South Qld: Ground and aerial application: Commence spraying at the start of the summer rainy season. Apply a maximum of 5 Group 3 Fungicides sprays per season. Apply a maximum of 2 consecutive sprays of Bulls-eye® at 21 to 28 day intervals before applying further treatments of a protectant fungicide from a different activity group. NT, Nth Qld: Ground and Aerial Applications: Commence spraying at the start of the wet season. Apply a regular schedule of protectant sprays. When the disease potential is high, apply a maximum of 2 consecutive sprays of Bulls-eye® at 14 to 21 day intervals before applying further treatments of a fungicide from a different activity group. Apply a maximum of 6 Group 3 sprays per season. Continue with treatments using an alternative protectant fungicide for the remainder of the season. Use the lower rate of oil in north Qld. DO NOT apply during July, August, September and October. * See Special Warning – Bananas in COMPATIBILITY for further information.
Carrots	Leaf Blight (<i>Alternaria dauci</i> and <i>Cercospora carotae</i>)	All States	300 or 500 mL	7	This use is subject to a CropLife Australia Fungicide Resistance Management Strategy. Use in a protectant program commencing before disease occurs. Limit the use of Group 3 Fungicide to periods when conditions favour disease development. Apply the 300 mL rate at 7 day intervals and the 500 mL rate at 10 to 14 day intervals. Use the 500 mL rate when conditions favour high disease pressure. Apply no more than 2 consecutive sprays of Bulls-eye® alone. Then apply a recommended non Group 3 Fungicide or tankmix Bulls-eye® with a recommended protectant fungicide for at least 1 application before continuing with Bulls-eye® alone. Good crop hygiene will aid disease control.

Macadamia Nuts	Husk Spot (<i>Pseudocercospora macadamiae</i>)	Qld, NSW, NT only	Dilute spraying 50mL/100 L water Concentrate spraying: Refer to the Application section	-	Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Use in a protectant fungicide program containing fungicides from different chemical groups. Commence fungicide applications at nut set (approximately late September) and continue the fungicide program until late December, with applications at 3 to 4 week intervals. Apply a maximum of 2 applications of Bulls-eye® per season. Ensure thorough spray coverage. Add Agral Spray Adjuvant at a rate of 10 mL/100 L.
Potatoes	Target Spot/Early Blight (<i>Alternaria solani</i>)	All States	300 or 500 mL	7	This use is subject to a CropLife Australia Fungicide Resistance Management Strategy. Use in a protectant program commencing when rows meet and before disease occurs. Limit the use of Group 3 Fungicides to periods when conditions favour disease development. Apply the 300 mL rate at 7 day intervals and the 500 mL rate at 10 to 14 day intervals. Use the 500 mL rate when conditions favour high disease pressure. Apply no more than 2 consecutive sprays of Bulls-eye® alone. Then apply a recommended non Group 3 Fungicide or tank mix Bulls-eye® with a recommended protectant fungicide for at least 1 application before continuing with Bulls-eye® alone. Good crop hygiene will aid disease control. Bulls-eye® does not control Late Blight (<i>Phytophthora infestans</i>). If this disease is a problem, mix Bulls-eye® with an appropriate fungicide for control of this disease and observe the spray interval for that product.
Tomatoes	Target Spot <i>Alternaria solani</i>	All States	300 or 500 mL	3	This is subject to a CropLife Australia Fungicide Resistance Management Strategy. Use in a protectant program commencing soon after transplanting. Limit the use of Group 3 Fungicides to periods when conditions favour disease development. Apply the 300 mL rate at 7 day intervals and the 500 mL rate at 10 day intervals. Apply no more than 2 consecutive sprays of Bulls-eye® alone. Then apply a recommended non Group 3 Fungicide or tank mix Bulls-eye® with a recommended protectant fungicide for at least 1 application before continuing with Bulls-eye® alone. DO NOT apply more than 6 Bulls-eye® (or other Group 3 Fungicide) sprays in a season. Good crop hygiene will aid disease control.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

BANANAS: **DO NOT** HARVEST FOR 1 DAY AFTER APPLICATION.

POTATOES, CARROTS: **DO NOT** HARVEST FOR 7 DAYS AFTER APPLICATION.

TOMATOES: **DO NOT** HARVEST FOR 3 DAYS AFTER APPLICATION.

AVOCADOES, MACADAMIAS: NO WITHHOLDING PERIOD WHEN USED AS DIRECTED.

NOTICE TO BUYER

To the extent permitted by the Competition and Consumer Act (2010) or any relevant legislation of any State or Territory (the "Legislation") all conditions and warranties and statutory or other rights of action, whether arising in contract or tort or whether due to the negligence of FMC or Seller, which buyer or any other user may have against FMC or Seller are hereby excluded provided however that any rights of the buyer pursuant to non-excludable conditions or warranties of the Legislation are expressly preserved. FMC hereby gives notice to buyer and other users that to the extent permitted by the Legislation it will not accept responsibility for any indirect or consequential loss of whatsoever nature arising from the storage, handling or use of this Product. Where permitted by the Legislation FMC's liability shall in all circumstances be limited to the replacement of the product, or a refund of the purchase price paid therefor.

The Product must be used and applied strictly in accordance with the label instructions and other directions for use. It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of FMC or the Seller. Buyer accepts these risks.

IN A MEDICAL EMERGENCY CALL 1800 033 111

ALL HOURS

The information set forth herein is furnished free of charge and is based on technical data that FMC believes to be reliable. It is intended for use by persons having technical skill at their own discretion and risk. FMC makes no warranties, expressed or implied, and assumes no liability in connection with any of this information. Nothing herein is to be taken as license to operate under or a recommendation to infringe on any patents.

FMC and Bulls-eye® Systemic Fungicide are trademarks of FMC Corporation or an affiliate.

© 2021 FMC Corporation. All rights reserved.

*Not an FMC trademark.

APVMA Approval No: 61336/61895

FMC Australasia Pty Ltd.
Level 2, Building B12 Julius Ave
North Ryde NSW 2113
Phone: 1800 066 355
www.fmccrop.com.au

SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111

ALL HOURS – AUSTRALIA WIDE
