

# POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Cheminova

# Chlorsulfuron 750 WG

Herbicide

ACTIVE CONSTITUENT: 750 g/kg CHLORSULFURON

GROUP

B

HERBICIDE

A selective herbicide for the control of Annual (Wimmera) Ryegrass and certain broadleaved weeds in Wheat, Barley, Oats, Cereal Rye and Triticale

CHEMINOVA  
AUSTRALIA PTY LTD  
ABN 59 472 616 773  
ACN 110 199 169  
12 JULIUS AVE  
NORTH RYDE NSW 2113  
PH: 1800 624 597  
[www.cheminova.com.au](http://www.cheminova.com.au)

CONTENTS: 500 g-50 kg

## DIRECTIONS FOR USE

### RESTRAINTS:

DO NOT spray emerged crops if rain is expected within four hours.

After mixing in the tank, spray within 48 hours if this product is used by itself, or within 24 hours if mixed with another product.

DO NOT apply to plants suffering stress.

## METHOD OF USE - PRE-SOWING INCORPORATED BY SOWING

### CROPS: WHEAT AND TRITICALE ONLY

Weeds Controlled	State(s)	Rate g/ha			Critical Comments
		Soil Type			
		Light to Medium Soils	Heavy Soils		
		Soil pH			
		Less than 7	7.0 - 8.5	8.5 or less	
Annual (Wimmera) Ryegrass <i>Lolium rigidum</i>	NSW, Vic, SA, WA only	20	15 or 20*	20	* Use the higher rate when paddock history suggests a high weed population can be expected. NOTE: Refer to General Instructions for optimum application timing and conditions.
Weeds Controlled	State(s)	Rate g/ha		Critical Comments	
African Turnip Weed <i>Sisymbrium thellungii</i>	Qld, NSW, ACT only	20			
Amsinckia/Yellow Burrweed <i>Amsinckia</i> spp.	NSW, ACT, Vic, SA, WA only	15			
Annual Phalaris <i>Phalaris paradoxa</i> , <i>Phalaris minor</i>	NSW, ACT only	20 + 1 L/ha Trifluralin	If possible, spray and incorporate into the soil in one operation. If this is not possible, incorporation should take place within four (4) hours of spraying. Delay may cause inferior weed control. Use only trifluralin products with an active level of 400 g/L.		
Barley grass <i>Hordeum leporinum</i>	NSW, ACT, Tas only				
Silvergrass <i>Vulpia</i> spp.	Tas only				
Ball Mustard <i>Neslia puniculata</i>	SA only	15			
Black Bindweed/Climbing Buckwheat <i>Fallopia convolvulus</i>	Qld only	20		Apply to dry soil before the sowing rain. Mechanical incorporation before the sowing rains is not necessary.	
Brome grass <i>Bromus</i> spp. (Suppression only)	NSW, ACT, Vic, Tas, SA, WA only	20		Gives suppression only if populations are 20 plants/m <sup>2</sup> or less.	
Cape Tulip <i>Homeria</i> spp.	WA only	20			
Capeweed <i>Arctotheca calendula</i>	NSW, ACT, Vic, Tas, SA, WA only	20		On acid soils pH 5.5 or less, this product will give a shorter period of control in wet years.	
Charlock <i>Sinapis arvensis</i>	Vic, Tas, SA only	15			
Common Iceplant <i>Mesembryanthemum crystallinum</i>	SA only	15			
Corn Gromwell/Sheepweed, White Ironweed <i>Buglossoides arvensis</i>	Qld, , ACT, Vic, SA, WA only	20			
Deadnettle <i>Lamium amplexicaule</i>	All States	15 or 20		Use the higher rate when paddock history suggests a high weed population can be expected.	
Docks <i>Rumex</i> spp.	NSW, ACT, Vic, Tas, SA, WA, only	20			
Fat-hen <i>Chenopodium album</i>	NSW, ACT, Tas only	20			
Fumitory <i>Fumaria</i> spp.	NSW, ACT, Vic, Tas, SA, WA, only	15 or 20		Use the higher rate when paddock history suggests a high weed population can be expected.	
Guildford grass/Onion grass <i>Romulea rosea</i>	WA only	15			
Indian Hedge Mustard <i>Sisymbrium orientale</i>	All States	15			
King Island Melilot <i>Melilotus indicus</i>	Vic, SA only	15			
Lincoln weed <i>Diplotaxis tenuifolia</i>	SA only	15			
Loosestrife	Vic only	15			
Mintweed <i>Salvia reflexa</i>	Qld, NSW, ACT, WA, only	20			
Mouse-ear Chickweed <i>Cerastium</i> spp.	NSW, ACT, Vic, Tas, SA, WA only	15			
New Zealand Spinach <i>Tetragonia tetragonoides</i>	Qld only	20			
Paradoxa grass <i>Phalaris paradoxa</i>	Nth NSW	20		Apply to dry soil before the sowing rain. Mechanical	

Weeds Controlled	State(s)	Rate g/ha	Critical Comments
	(Soil pH >7.5) and Qld only		incorporation before the sowing rains is not necessary.
Paterson's Curse/Salvation Jane <i>Echium plantagineum</i>	NSW, ACT, Vic, Tas, SA, WA, only	15	
Pimpernels <i>Anagallis arvensis</i>	NSW, ACT, Vic, Tas, SA, only		
Prickly Lettuce/Whip Thistle <i>Lactuca serriola</i>	Vic, SA only	20	
Rough Poppy <i>Papaver hybridum</i>	NSW, ACT, Tas, SA, WA only	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
Saffron Thistle <i>Carthamus lanatus</i> (Suppression only)	Qld, NSW, ACT, Vic, Tas, SA only	20	
Saltbush <i>Atriplex muelleri</i>	Qld, NSW, ACT, only		
Shepherd's Purse <i>Capsella bursa-pastoris</i>	NSW, ACT, Vic, Tas, SA, WA only	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
Slender Celery <i>Apium leptophyllum</i>	Qld, NSW, ACT only	20	
Slender Thistle <i>Carduus tenuiflorus</i>	Tas only		
Soursob <i>Oxalis pes-caprae</i>	NSW, ACT, Vic, SA only	15	Apply only to soils of pH 7.5 or above. Apply after majority of Soursobs have emerged and leave soil undisturbed for 1-4 weeks prior to cultivation or sowing. The most effective and reliable control is achieved with early post-emergence applications (EPE) after crop and weed emergence.
Spear Thistle <i>Cirsium vulgare</i>	Tas only	20	
Stemless Thistle <i>Onopordum acaulon</i>	SA only	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
Storksbill/Wild Geranium <i>Erodium</i> spp.	Vic, Tas, SA, WA only	15	
Three-Cornered Jack(s)/Doublegee/Spiny Emex <i>Emex australis</i>	NSW, ACT, Vic, SA, WA only	20	
Tree Hogweed <i>Polygonum patulum</i>	Vic, SA only		
Turnip weed <i>Rapistrum rugosum</i>	Qld and SA only	15	
Wireweed/Hogweed <i>Polygonum aviculare</i>	All States	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
Wild Turnip <i>Brassica tournefortii</i>	NSW, ACT, Vic, Tas, SA, WA only	15	

## METHOD OF USE - POST CROP AND WEED EMERGENCE

### CROP: WHEAT, BARLEY, OATS, TRITICALE AND CEREAL RYE ONLY

Weeds Controlled	State(s)	Rate g/ha			Critical Comments
		Soil Type			
		Light to Medium Soils	Heavy Soils		
		Soil pH			
		Less than 7	7.0 - 8.5	8.5 or less	
Annual (Wimmera) Ryegrass <i>Lolium rigidum</i>	NSW, Vic, SA, WA only	20 or 25*	15 or 20*	20 or 25*	* Use the higher rate under heavy weed pressure. Apply no later than the 3 leaf stage of Annual Ryegrass. * Application of This product to Annual Ryegrass 2 leaf or greater with water volumes less than 50 L/ha may result in reduced efficacy.
Weeds Controlled	State(s)	Rate g/ha	Critical Comments		
African Turnip weed <i>Sisymbrium thellungii</i>	Qld, NSW, ACT, only	20	Apply at cotyledon to 4 leaf stage.		
Amsinckia/Yellow Burrweed <i>Amsinckia</i> spp.	NSW, ACT, Vic, SA, WA only	15			
Ball Mustard <i>Neslia puniculata</i>	SA only				
Bifora/Carrot weed <i>Cotula australis</i>		25	Apply at cotyledon to 2 leaf stage of weed.		
Black Bindweed/Climbing Buckwheat <i>Fallopia convolvulus</i>	Qld, NSW, ACT, only	20			
Cape Tulip <i>Homeria</i> spp.	WA only				
Charlock <i>Sinapis arvensis</i>	NSW, ACT, Vic, Tas, SA only	15			
Corn Gromwell/Sheepweed/ White Ironweed <i>Buglossoides arvensis</i>	NSW, ACT, Vic, SA, WA only	20	Apply at cotyledon to 2 leaf stage. If applied at a later stage only suppression will occur.		

Weeds Controlled	State(s)	Rate g/ha	Critical Comments
Deadnettle <i>Lamium amplexicaule</i>	Qld, NSW, ACT, Vic, Tas, SA only	15 or 20	Use the higher rate under heavy weed pressure.
Docks <i>Rumex</i> spp.	Vic, Tas, SA, WA, only	15	
Fat-hen <i>Chenopodium album</i>	NSW, ACT, Tas only	20	
Fumitory, Denseflower <i>Fumaria densiflora</i>	NSW, ACT, Vic, Tas, SA, WA, only	20	Apply at cotyledon to 2 leaf stage.
Guildford grass/Onion grass <i>Romulea rosea</i>	WA only	15	
Hoary Cress <i>Cardaria draba</i>	Vic, Tas, SA, only	20	Apply when plants are fully emerged.
Lincoln weed <i>Diplotaxis tenuifolia</i>	SA only	20	
Matricaria <i>Matricaria matricarioides</i>	Tas, WA only		
Mintweed <i>Salvia reflexa</i>	Qld, NSW, ACT only		Apply at cotyledon to 4 leaf stage.
Mouse-ear Chickweed <i>Cerastium</i> spp.	NSW, ACT, Vic, Tas, SA, WA, only	15	
Mustards <i>Sisymbrium</i> spp.	All States		
New Zealand Spinach <i>Tetragonia tetragonoides</i>	Qld only	20	
Paterson's Curse/Salvation Jane <i>Echium plantagineum</i>	NSW, ACT, Vic, Tas, SA, WA, only	15	
Pimpernels <i>Anagallis arvensis</i>	NSW, ACT, Vic, Tas, SA only		
Prickly Lettuce/Whip Thistle <i>Lactuca serriola</i>	Vic, Tas only	20	
Rough Poppy <i>Papaver hybridum</i>	NSW, ACT, Tas, SA, WA, only		
Saltbush <i>Atriplex muelleri</i>	Qld, NSW, ACT only		Apply at cotyledon to 4 leaf stage.
Shepherd's Purse <i>Capsella bursa-pastoris</i>	NSW, NSW, ACT, Vic, Tas, SA, WA only		
Slender Celery <i>Apium leptophyllum</i>	Qld, NSW, ACT only		Apply at cotyledon to 4 leaf stage.
Soursob <i>Oxalis pes-caprae</i>	NSW, ACT, Vic, SA, WA only		Apply when the majority of Soursobs have emerged.
Spear Thistle <i>Cirsium vulgare</i>	Tas only		
Stagger weed <i>Stachys arvensis</i>	Qld, NSW, ACT, Tas, WA only		
Stemless Thistle <i>Onopordum acaulon</i>	Vic only	25	
Storksbill/Wild Geranium <i>Erodium</i> spp.	Vic, Tas, SA, WA only	15	
Tree Hogweed <i>Polygonum patulum</i>	Vic only	20	
Turnip Weed <i>Rapistrum rugosum</i>	Qld, NSW, ACT, SA only	15	
Wild Radish <i>Raphanus raphanistrum</i>	All States	15 or 20	Use the higher rate under heavy weed pressure. A follow-up spray with a suitable herbicide may be necessary to control subsequent germinations.
Wild Turnip <i>Brassica tournefortii</i>	NSW, ACT, Vic, Tas, SA, WA only	15	
Wireweed/Hogweed <i>Polygonum aviculare</i>	All States	20	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED**

#### GENERAL INSTRUCTIONS

This product is a selective herbicide designed to control certain weeds in wheat, triticale, barley, oats and cereal rye. This product is suitable as a pre-sowing treatment for wheat and triticale, and as a post-sowing treatment for wheat, triticale, barley, oats and cereal rye. When used on emerged weeds, the product is absorbed by foliage and green stems and moves into the root system.

Prior to using this product, careful consideration should be given to soil pH. As soil pH increases, rate of breakdown decreases. This product should not be used on soil pH 8.6 or higher as extended soil residual activity could adversely affect crop rotation options beyond normal intervals.

This product should not be used on soils with a pH of 8.6 or higher as soil residual activity could adversely affect following crops and crop rotation intervals may be extended beyond normal intervals.

Crops other than wheat, barley, oats, triticale and cereal rye can be extremely sensitive to low concentrations of this product in the soil. See Crop Rotation Recommendations.

Best weed control is obtained when rainfall or sprinkler irrigation wets the soil to a depth of 5 to 7.5 cm within 4 weeks of application.

### PRE-SOWING INCORPORATED BY SOWING:

WA only - Avoid applying to dry sandy soils as rapid leaching may occur with early season rains.

SA only - Before using rates greater than 15 g/ha on light to medium soils pH 7 to 8.5, seek further advice.

Conventionally Sown Crops - on soils less than 7, apply a spray just before sowing or in conjunction with the sowing operation. On soils of pH of 7 or greater it is not critical to time the spray just before sowing. Spray onto a non-ridged surface free of large clods. Use low profile 10 cm combine points for sowing. Sow at speeds of 10 kph or greater. Use light covering harrows at sowing. If applied to dry soil and sowing is to be delayed, incorporate immediately after spraying to prevent loss by wind erosion.

Direct Drilled Crops - apply tank mixed with either Spray.Seed<sup>^</sup> or glyphosate in accordance with manufacturer's label recommendations.

### POST CROP AND WEED EMERGENCE:

Where treatment is delayed or where weeds are not actively growing due to adverse conditions results may be slow to appear and weeds may be only stunted or suppressed.

Wheat, Triticale and Cereal Rye - apply after crop emergence when weeds are small and actively growing (**Annual Ryegrass no more than 3 leaves, Broadleaved weeds no more than 5 cm in height or diameter** (for Black Bindweed refer to specific recommendations)).

Barley and Oats - apply between 2 leaf stage of crop (3 leaf stage - SA only) and early tillering when weeds are small and actively growing (**Annual Ryegrass no more than 3 leaves, Broadleaved weeds no more than 5 cm in height or diameter** (for Black Bindweed refer to specific recommendations)).

### RESISTANT WEEDS WARNING

GROUP	<b>B</b>	HERBICIDE
-------	----------	-----------

Cheminova Clorsulfuron 750WG Herbicide is a member of the sulfonylurea group of herbicides. Cheminova Clorsulfuron 750WG Herbicide has the inhibitor of the enzyme acetolactate synthase (ALS) mode of action. For weed resistance management, Cheminova Clorsulfuron 750WG Herbicide is a Group B herbicide.

Some naturally occurring weed biotypes resistant to the product and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Cheminova Clorsulfuron 750WG Herbicide or any other Group B herbicides.

Annual Ryegrass biotypes resistant to diclofop-methyl and other "grass specific" herbicides are often also resistant to Cheminova Clorsulfuron 750WG Herbicide. Before using Cheminova Clorsulfuron 750WG Herbicide on a population resistant to "grass specific" herbicides, have a resistance test conducted to ensure that it is still susceptible to Cheminova Clorsulfuron 750WG Herbicide.

Since the occurrence of resistant weeds is difficult to detect prior to use, Cheminova Australia Pty Ltd accepts no liability for any losses that may result from the failure of Cheminova Clorsulfuron 750WG Herbicide to control resistant weeds.

To prevent, or at least minimise the risk of resistant weeds occurring, use Cheminova Clorsulfuron 750WG Herbicide in tank mixes (if appropriate) and/or rotations with herbicides having different modes of action effective on the same weed species.

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be taken to prevent seed set of these survivors.

DO NOT make more than one application of an ALS inhibitor herbicide to a crop, either pre-sowing incorporated by sowing or post crop and weed emergence.

If the user suspects that an ALS inhibitor-resistant weed is present, Cheminova Clorsulfuron 750WG Herbicide or other Group B herbicides should not be used.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

### GRAZING ADVICE

Avoid grazing treated areas within 24 hours of application to optimise weed control.

A nil withholding period is applicable for grazing Cheminova Clorsulfuron 750WG Herbicide treated areas (when used as directed on this label).

### CROP SAFETY

DO NOT use this product for:

- crops other than cereals.
- cereals irrigated by furrows or flooding.
- winter cereals undersown with legume pasture crops.

- weed control where crops are under stress. Damage can occur where crops are stressed due to conditions such as excessive soil alkalinity or acidity, poor nutrient status, disease, nematode or insect infestation, adverse weather conditions, drought or waterlogging. If crops become stressed after spraying, they may turn yellow or become retarded, but usually they will recover with no reduction in yield.

**Wheat:**

DO NOT use this product for:

- wheat varieties Cranbrook, or Miling.
- the wheat variety Vulcan if on acid soils and under stress conditions caused by waterlogging, frost, aluminium or manganese toxicity; reduced yields may result.
- pre-sowing treatment of weeds in wheat varieties Avocet and Durati (OK for post-emergent use).
- pre-sowing treatment of weeds in wheat variety Banks if soil pH is 5.5 or less (OK for post-emergent use).

**Barley and Oats:**

DO NOT use this product for:

- application before the crop has reached the 2-leaf stage (3-leaf stage in SA).
- Stirling barley.
- barley under waterlogged conditions (yield may be reduced).

The application of other sulfonylurea herbicides following this product is not recommended.

**CROP ROTATION RECOMMENDATIONS**

Land previously treated with this product should not be rotated to other crops other than those listed in the following tables. Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas. The treated areas may be replanted to any of the specified crops after the interval indicated in the following tables:-

**NB - THE TABLE BELOW APPLIES TO ALL STATES**

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)						
SOIL pH*	0	3	6	9	12	18
6.5 or less	Triticale Wheat	Cereal Rye	Oats	Barley	Subterranean Clover** Faba Beans Field Pea Linseed Lucerne Lupins Medics ** Canola/Rapeseed Safflower	Maize Sorghum Soybeans Sunflower

**NB - THE TABLES BELOW APPLY TO Qld, SA, WA & Tas ONLY**

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)						
RAINFALL REQUIREMENT	0	3	9	15	18	22
	Minimum 700 mm					
Soil pH* 6.6-7.5	Triticale Wheat	Cereal Rye	Barley Oats	Japanese Millet Maize Panicum Millet Sorghum Sunflower White French Millet	Cotton Soybeans	Faba Beans Field Pea Linseed Medics** Canola/ Rapeseed Safflower Subterranean Clover**

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)				
RAINFALL REQUIREMENT	0	15	18	24 months or longer
	Minimum of 700 mm			
Soil pH* 7.6-8.5	Triticale Wheat	Japanese Millet Maize Panicum Millet Sorghum Sunflowers White French Millet	Barley Oats Cereal Rye	Rotate to crops other than Cereals (such as listed above) only if field test strip of the planned rotational crop has been successfully grown through to maturity in the previous season.
8.6 and above	This product is not recommended for use on soils of pH 8.6 and above.			

**NB - THE TABLES BELOW APPLY TO NSW & Vic ONLY**

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)					
SOIL pH*	0	3	9	22	26
6.6 - 7.5	Triticale Wheat	Cereal Rye	Barley Oats	Subterranean Clover** Faba Beans Field Peas Linseed Lucerne Lupins Medics ** Canola/Rapeseed Safflower	Maize Sorghum Soybeans Sunflower

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)			
SOIL pH*	0	18	24 months or longer
7.6 - 8.5			Rotate to crops other than Cereals (such as listed above) only if field test strip of the planned rotational crop has been successfully grown through to maturity in the previous season.
8.6 and above	This product is not recommended for use on soils of pH 8.6 and above.		

\* Soil pH is to be determined by Laboratory analysis using the 1:5 soil:water suspension method.

\*\* Includes natural regeneration of Subterranean clover and Medics.

- Land previously treated with this product should not be rotated to crops other than those listed in the above table.
- Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas.

**SPRAY PREPARATION**

This product is a water dispersible granule.

1. Fill tank partially with water and engage full agitation.
2. Add the required amount. (N.B. The measuring flask provided is graduated in grams of Cheminova Chlorsulfuron 750WG Herbicide only. DO NOT use for measuring of other materials.)
3. Top up with water to the required volume.
4. Companion products: If applying this product with another product ensure this product has completely dissolved before adding the companion product.
5. Cheminova Clorsulfuron 750WG Herbicide must be kept in suspension at all times by continuous agitation. Where prepared spray mixes have been allowed to stand, thoroughly re-agitate before using.

**USE OF SURFACTANT/WETTING AGENT**

For post-emergence application, always add a non-ionic surfactant (1000 gac/L) at 100 mL/100 L of final spray volume (0.1% volume/volume). The use of spraying oils is not recommended.

NOTE: DO NOT add a surfactant/wetting agent when this product is tank mixed with another product which already has a surfactant/wetting agent in the formulation.

**GROUND SPRAYING EQUIPMENT**

Use a boom spray properly calibrated to a constant speed and rate of delivery to ensure thorough coverage and a uniform spray pattern. Avoid overlapping and shut off spray booms while starting, turning, slowing or stopping as injury to the crop may result. Apply a minimum of 30 L prepared spray/ha.

**AERIAL APPLICATION**

Apply a minimum of 20 L/ha water. Avoid spraying in still conditions and in winds likely to cause drift onto adjacent sensitive crops. Avoid spraying where drift can go onto areas likely to be sown to sensitive crops - see Crop Rotation Recommendations. Turn off spray boom whilst passing over creeks and dams.

**SPRAYER CLEANUP**

It is essential that the sprayer be properly cleaned after using this product prevent injury to crops other than wheat, triticale, barley, oats, or cereal rye. All traces of Cheminova Chlorsulfuron 750WG Herbicide should be removed from equipment using the following procedure:

1. Drain tank, then flush tank, boom and hoses with clean water for a minimum of 10 minutes.
2. Fill the tank with clean water then add 300 mL household chlorine bleach (containing 4% chlorine) per 100 L of water. Flush through boom and hoses then allow to sit for 15 minutes with agitation engaged, then drain.
3. Repeat Step 2.

4. Nozzles and screens should be removed and cleaned separately. To remove traces of chlorine bleach, rinse the tank thoroughly with clean water and flush through hoses and boom.

**CAUTION:** DO NOT use chlorine bleach with ammonia. All traces of liquid fertiliser containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odour which can cause eye, nose, throat and lung irritation. DO NOT clean equipment in an enclosed area.

#### **COMPATIBILITY**

This product is compatible with glyphosate and paraquat. This product does not control wild oats, however it is compatible with wild oat herbicides triallate, Mataven<sup>^</sup> and Foxtrot<sup>®</sup>. It is also compatible with bromoxynil, MCPA (and bromoxynil/MCPA mixtures), 2,4-D amine and 2,4-D ester, clopyralid, Nimble<sup>®</sup> and Jester<sup>®</sup>. This product is also compatible with trifluralin and the insecticides LeMat<sup>^</sup>, Danadim<sup>®</sup> deltamethrin and Cyren<sup>®</sup>.

#### **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply or drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

#### **STORAGE AND DISPOSAL**

Store in the closed, original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

#### **SAFETY DIRECTIONS**

Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

#### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 131126)

#### **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet available from [www.cheminova.com.au](http://www.cheminova.com.au)

#### **WARNING**

This product must be used, handled and stored strictly as directed in accordance with this label, packaging and other reference material ("Directions"). To the extent permitted by law Cheminova Australia Pty Ltd and its related companies will have no liability for any injury, loss or damage arising from a failure to follow the Directions.

APVMA Approval No: 59542/62156

Batch No:

Date of Manufacture:

140213-T

Drummuster logo (relevant pack sizes)

<sup>®</sup> Registered trademark of Cheminova A/S    <sup>^</sup> Trademark

<p><b>FOR 24 HOUR SPECIALIST ADVICE IN EMERGENCY ONLY PHONE 1800 033 111</b></p>
--