

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Cheminova

MCPA 750

Herbicide

ACTIVE CONSTITUENT: 750 g/L MCPA
present as the DIMETHYLAMINE SALT

GROUP

I

HERBICIDE

For the selective control of broadleaf weeds in cereals, linseed pastures, rice, sugarcane and turf as per the Directions for Use Table

CHEMINOVA AUSTRALIA PTY LTD

ABN 59 472 616 773

ACN 110 199 169

12 JULIUS AVE

NORTH RYDE NSW 2113

PH: 1800 624 597

www.cheminova.com.au

CONTENTS: 10-1000 L

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply if rain is likely within 6 hours.

1. CEREALS

| CROP | WEEDS CONTROLLED | STATE | RATE/ha | CRITICAL COMMENTS |
|--|--|------------------------------------|---|---|
| Wheat, Barley, Oats, Triticale | Refer to Weed Table | SA, Tas only | 460 mL-1.45 L | Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing. |
| | | Qld only | | Apply from early tillering (4-5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem). |
| | | NSW, ACT only | | Apply 460 mL rate after crop has reached 5 leaf stage and weeds marked # in weed table have 3-6 leaves. Higher rates after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller. |
| | | Vic only | 460-960 mL | Apply 460 mL rate after crop has reached 2-3 leaf stage and weeds marked # in weed table have 2-4 leaves. Higher rate after the crop plants have 5 leaves to fully tillered. |
| | | WA only | 600 mL-1.35 L | Apply to crop from 5 leaf to Flag leaf emergence (Zadoks 15-37). |
| Soldier Thistle | Skeleton Weed | SA only | 460 mL plus 200 mL Cheminova Clopyralid | Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing. |
| | | | 660 mL plus 500 mL Sparta 300 | Apply to rosettes from 5-15 cm diameter when the crop is between the 5 leaf and late tillering growth stages. It is essential that all the Skeleton Weed is emerged and minimum of 5 cm diameter which is usually not before August. This rate will provide control of Skeleton Weed until harvest and will substantially reduce seed numbers the following season. |
| Wheat, Barley, Triticale, Cereal Rye | Capeweed, Doublegee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish | WA only | 270 mL plus 350 mL Flowable Diuron | Crop: 3-4 leaf stage (Zadoks 13-14). |
| | | | 330 mL plus 400 mL Flowable Diuron | Crop: 4-5 leaf stage (Zadoks 14-15). |
| | | | | Apply by aircraft or boom with 20-100 L/ha water, when most weeds have germinated and are in 1-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. DO NOT apply to undersown medic. |
| | Amsinckia (Yellow Burr), Capeweed, Charlock, Deadnettle, Doublegee (Spiny Emex, Three Corner Jack), <i>Erodium</i> spp., Mustard, Turnip Weed, White Ironweed (Corn Gromwell, Sheepweed), Wild Radish, Wild Turnip | NSW, ACT, Vic only | 330 mL plus 500 mL Flowable Diuron | Apply by boom spray when the crop is at the 3-5 leaf stage (NSW and SA only) and 5 leaf stage (Vic only). Weeds must be in the 2-5 leaf stage. DO NOT apply to undersown medic or lucerne. DO NOT use on Kamilaroi, Kite, Olympic or Shortim Wheat. DO NOT use on very light sandy soils. TANK MIXTURES: Read and follow all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use re restrictions and safety directions for the tank mix products. |
| Capeweed, Mustard, Three Cornered Jack, Turnip Weed, Wild Radish, Wild Turnip, Wards Weed, Common Ice Plant, Pimpernel | SA only | 230 mL plus 500 mL Flowable Diuron | | |

2. OTHER FIELD CROPS

| CROP | WEEDS CONTROLLED | STATE | RATE/ha | CRITICAL COMMENTS | |
|------------|-------------------------|-----------------------------|---------------|--|---|
| Field Peas | Prickly Lettuce | NSW, ACT, Vic, Tas, SA only | 4 leaf | 80 mL plus 125 mL Pelican | Field Peas: Apply early post-emergence after the third node stage and before the start of flowering. Warning: Field peas grown on high pH soils in the presence of free lime may be less tolerant to MCPA 750. APPLICATION AND WEED CONTROL Apply when weeds are actively growing. For optimum results apply 4 to 6 weeks post-sowing. Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: <ul style="list-style-type: none"> • high crop and weed density; • late season germinations; • abnormal weed growth (including early flowering); • higher rates of product (up to the maximum rate of application specified for that weed) may be required. MCPA 750 will not effectively control: <ul style="list-style-type: none"> • regrowth of suppressed weeds; • transplanted weeds; • regrowth from rhizomes or roots; • weeds growing under stress from previous herbicide applications. |
| | | | 6 leaf | | |
| | Charlock (Wild mustard) | | 4 leaf | 80 mL plus 125 mL Pelican | |
| | | | 6 leaf | 100 mL plus 150 mL Pelican | |
| Flax | Refer to Weed Table | Vic, Tas only | 325 mL | Apply to crop when 8-20 cm high. | |
| Linseed | Refer to Weed Table | Qld only | 460 mL-1.45 L | Apply when crop is 10-15 cm tall with at least 170 L/ha of water. DO | |

| CROP | WEEDS CONTROLLED | STATE | RATE/ha | CRITICAL COMMENTS |
|------------------------|---|-----------------------------|---------------|---|
| | | NSW, ACT, Vic, SA, Tas only | 460-730 mL | NOT spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. DO NOT apply by aircraft. |
| Oilseed Poppies | Fat Hen, Spear Thistle | Tas only | 1 L | Use only in consultation with the poppy contracting company. Apply at least one week after petal fall after poppy seed has been set. Apply by aircraft in 30-40 L of water/ha. This treatment will control late germinating weeds if they are green and at the bud or early flowering stage. Spraying at this time will twist weeds below the crop canopy. DO NOT apply in horticultural areas where there is a risk of spray or vapour drift over susceptible crops. |
| Rice | Dirty Dora, Starfruit, Sedges, Buddah (Butter) Pea | NSW only | 485 to 970 mL | Apply by aircraft at early/mid tillering stage of rice but not later than panicle initiation. Use 484 mL/ha at 35 days after aerial sowing (approx. 2 litres per rice plant). Use 970 mL/ha at 45 days or more after cereal sowing (3 tillers or more per rice plant). The crop should be sprayed before flowering of the weeds. Water Management: Before herbicide application, lower water level in the bays to at least two thirds of the weed growth for direct contact with MCPA 750. Weeds covered by water will not be controlled. Raise water depth to normal levels 24 hours following spraying. |
| Sugarcane | Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory | Qld only | 930 mL | Post-Emergent: Apply as directed inter-row spray. DO NOT exceed rates with phenoxy sensitive varieties. |
| | As above plus Fat Hen, Noogoora Burr | | 1.45 L | |

2. PASTURES, SEED CROPS & TURF

| SITUATION & CROP | WEEDS CONTROLLED | STATE | RATE/ha | CRITICAL COMMENTS |
|---|--|----------------------------|--|---|
| Grass Pastures | Refer to Weed Table | All States | 460 mL-2.7 L | Apply to established grass based pastures. Damage may occur to Legumes if present. See "Legume Tolerance" below. Note: Capeweed, Paterson's Curse and Variegated Thistle may be poisonous to stock after spraying. |
| | Soldier Thistle | SA only | 460 mL-1.45 L | Use low rate on seedlings with no more than 2 true leaves. Use 100 litres of water for best results. If clover present do not exceed 575 mL. |
| | Seedling and established Horehound | NSW, ACT only | 2 L | Apply in Autumn when horehound plants are at the seedling stage. If Sub-clover or Medic plants are present they will be damaged by this application. |
| Grass Seed Crops | Refer to Weed Table | Qld, Vic, Tas, SA, WA only | 460 mL-2.7 L | Apply to established grass seed crops. |
| Pastures – Spray / Graze Technique | Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherds Purse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip | NSW, ACT, Vic only | 230-930 mL | Spray actively growing 6-8 week old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. Refer "Legume Tolerance" under General Instructions. Precaution: An increased quantity of poisonous plants may be eaten by stock using Spray/Graze, eg. Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. DO NOT graze horses or pigs on Paterson's Curse. Observe Withholding Period of at least 7 days after spraying. |
| | Refer to Weed Table | WA only | 610 mL-1.25 L | |
| | Capeweed, Dandelion, Seedling Dock, Three Cornered Jack, Geranium, Erodium, Mustard, Annual Thistles, Turnip Weed, Wild Radish, Wild Turnip | SA only | 330 mL | |
| | Soldier Thistle | | 395 | |
| Sub-clover seed crops | Refer to Weed Table | Vic, SA only | 460-960 mL | DO NOT exceed rate specified. Spray in Autumn and early Winter. |
| Turf | Refer to Weed Table | All States | 330 mL or 2.8 mL per 100 sq. metres | Newly laid – Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses. |
| | | | 930 mL-1.8L or 10-19 mL per 100 sq. metres | Established – Apply in high volume to actively growing weeds. DO NOT mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses. |

3. SPOT SPRAYING

| SITUATION & CROP | STATE | MIXING RATES/CRITICAL COMMENTS |
|----------------------|------------|---|
| High Volume Spraying | All States | Use 130 mL/100 L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. |
| Knapsack Application | | Use 270 mL/100 L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed. Use 13 mL/10 L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. Use 27 mL/10 L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed. |

4. NON CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY

| SITUATION & CROP | WEEDS CONTROLLED | STATE | RATE | CRITICAL COMMENTS |
|---|---------------------|------------|---------------------|--|
| Non Crop Areas, Fallow Land, Industrial & Commercial Areas, Rights of Way | Refer to Weed Table | All States | Refer to Weed Table | Apply at rates shown in Weed Table when weeds only are present and no crop or pasture is involved. |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

CEREALS, PASTURE: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED

OILSEED POPPIES: DO NOT Harvest for 5 weeks after application

WEED TABLE:

Where weeds are to be sprayed in a crop or pasture use only the rates given for particular crop or situation indicated under the Directions for Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

APPLICATION RATE, LITRES PER HECTARE

*#: refer to Critical Comments for Wheat, Barley, Oats, Triticale

| WEEDS CONTROLLED | Vic | NSW, ACT | SA | Tas | Qld | WA | CRITICAL COMMENTS |
|---------------------------|-------|-----------|------|-----------|-----------|-------|--|
| Ball Mustard | - | - | 1.15 | - | - | - | Apply at rosette stage. |
| Bathurst Burr | 1.35 | 0.66-1.35 | 1.35 | 1.35 | 1.35 | 1.35 | Spray young seedlings only. |
| Black Bindweed | - | 0.97-1.35 | - | - | 0.93-1.35 | - | Apply at young stage prior to flowering. |
| Californian Burr | - | 0.66-1.35 | - | 1.45 | - | - | Spray young seedlings only. |
| Cape Tulip | - | - | - | - | - | 1.35 | Apply just prior to flowering. Gives mild suppression only. |
| Capeweed | - | 1.45 | 1.45 | 1.45 | 1.45 | - | Apply at seedling stage. |
| Carrot Weed | 1.35 | - | - | - | 1.35 | - | Apply at seedling stage. |
| Charlock | 0.46 | 0.66 | 0.46 | 0.66-1.45 | 0.46 | - | Spray up to rosette stage. |
| Cutleaf Mignonette | - | - | 1.45 | - | - | - | Apply at young rosette stage. |
| Dandelion | 1.35 | 1.35 | 1.35 | 1.35 | 1.35 | 1.35 | Apply at young rosette stage. |
| Deadnettle | - | 1.45 | - | - | 1.45 | - | Moderately susceptible. Apply at seedling stage. |
| Docks | 0.96 | - | 1.45 | 1.45 | - | - | Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected. |
| Fat Hen | 1.35 | 0.66-1.35 | - | 1.35 | 1.35 | - | Spray up to rosette stage. |
| Fennel | 1.80 | 1.80 | 1.80 | 1.80 | 1.80 | - | Apply at young stage. |
| Field Bindweed | 1.35 | - | 1.35 | 1.35 | 1.35 | 1.35 | Apply at bud stage, repeating over several seasons. |
| Field Cress | - | - | - | 1.45 | - | - | Apply at young stage. |
| Flatweed or Catsear | - | - | - | 1.45 | - | 0.93 | Apply when weeds are 10-15 cm high. |
| Fumitory (red) | 0.93 | 0.93 | 0.93 | - | 0.93 | - | Apply at young stage prior to flowering. |
| Hoary Cress | 3 | - | 1.45 | - | 3 | - | Apply before flowering. |
| Horehound | - | 2 | 1.45 | - | - | - | Apply at rosette stage. |
| Hedge Mustard | 0.73# | - | 0.46 | 0.46 | 0.46 | 0.825 | Apply at rosette stage. |
| Hexham Scent or Melilotus | 0.96 | 0.80 | 0.93 | - | 0.93 | - | Apply at young seedling stage. |
| Lesser Swinecress | - | - | - | 1.45 | - | - | Apply at young stage. |
| Lincoln Weed | - | - | 0.46 | - | - | - | Apply at rosette stage. |
| London Rocket | - | - | - | - | - | 0.825 | |
| Lupins | - | 0.46-0.96 | - | - | - | 0.825 | Spray up to 10 cm high. |
| Mintweed | 1.35 | 1.35 | - | - | 1.35 | - | Spray young seedlings only. |
| Mustards | 0.73# | 0.66# | 0.73 | 0.73 | - | 0.825 | Spray up to rosette stage. |
| Noogoora Burr | 1.35 | 0.66-1.35 | 1.35 | - | 1.35 | 1.35 | Spray young seedlings only. |
| Opium Poppy | - | - | - | 1.45 | - | - | Apply at young stage. |
| Paterson's Curse | - | 0.66-0.96 | - | - | 2.7 | - | Apply early rosette stage. |
| Pimpernel | - | - | 0.46 | - | - | - | Apply at rosette stage. |
| Plantains | - | 1.35 | - | 1.35 | 1.35 | - | Apply at young stage. |
| Rapeseed | - | 0.66-0.96 | - | - | - | - | Spray up to rosette stage. |
| Rough Poppy | - | 0.46-0.96 | - | - | - | - | Spray up to rosette stage. |

| WEEDS CONTROLLED | Vic | NSW, ACT | SA | Tas | Qld | WA | CRITICAL COMMENTS |
|--------------------------|-----------|-----------|-----------|-----------|-----------|-----------|--|
| Safflower | - | 0.46-0.66 | - | - | - | - | Spray up to rosette stage. |
| Shepherd's Purse | - | - | - | 1.45 | - | - | Apply at rosette stage. |
| Skeleton Weed | - | 0.96-1.35 | - | - | - | - | Spray rosettes before aerial growth commences. |
| Sorrel | - | - | 1.45 | - | - | - | Apply to rosette stage. |
| Stinkwort | 1.35 | 0.66-0.96 | 1.35 | - | - | 1.35 | Apply when weeds are 10-15 cm high. |
| Sunflower (seedlings) | - | 0.69-1 | - | - | - | - | Spray up to rosette stage. |
| Thistles: - Artichoke | 1.05 | - | 1.45 | - | 1.35 | 0.46 | Apply at rosette stage. |
| - Nodding | - | 1.65 | - | - | - | - | Spray early rosettes. Repeat spray required in following Spring or Autumn. |
| - Saffron | 0.96-1.65 | 0.66-1.35 | 0.93-1.65 | 0.93-1.65 | 0.93-1.65 | 0.96-1.65 | Spray up to rosette stage (September WA). |
| - Scotch or Cotton | 1.45 | - | 0.46-0.93 | 1.45 | - | - | Apply at seedling stage. |
| - Slender | 2 | 0.66-1.35 | 0.66 | 2 | - | 2 | Spray young seedlings only. |
| - Soldier | 0.73 | - | 0.73 | - | - | - | Spray young seedlings. Use higher rate for larger weeds. |
| - Spear | - | 0.96-1.35 | - | 1.45 | - | - | Apply at rosette stage. |
| - Star | - | - | 730-0.93 | - | - | - | Apply at seedling stage. Use higher rate for larger weeds. |
| - Variegated | 0.96 | 0.66-1.35 | 0.46-0.93 | 0.96 | 0.96 | 0.96 | Spray young seedlings only. Use 1.35 L/ha (SA) for well developed weeds. |
| Turnip Weed | 0.73# | 0.66# | 0.66 | - | - | - | Spray up to rosette stage. |
| Wild Artichoke | - | - | 1.45 | - | - | - | Apply at rosette stage. |
| Wild Radish | 0.93 | 0.66# | 0.93 | 1.35 | 0.46-0.73 | 0.80 | Spray up to rosette stage. |
| Wild Turnip | 0.73# | 0.66# | 0.46 | 1.35 | 0.46-0.73 | 0.60 | Spray up to rosette stage. |

GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application.

DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

APPLICATION/MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

Boom Spraying: Use 30-120 litres water per hectare.

Use minimum of 170 litres water per hectare for LINSEED.

Aerial Spraying: Use 10-90 litres water per hectare.

RESISTANT WEEDS WARNING

| GROUP | I | HERBICIDE |
|-------|---|-----------|
|-------|---|-----------|

Cheminova MCPA 750 Herbicide is a member of the Phenoxy group of herbicides. Cheminova MCPA 750 has the Disruptors of plant cell growth mode of action. For weed resistance management Cheminova MCPA 750 is a Group I herbicide. Some naturally-occurring weed biotypes resistant to Cheminova MCPA 750 and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Cheminova MCPA 750 or other disruptors of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Cheminova Australia Pty Ltd accepts no liability for any losses that may result from the failure of Cheminova MCPA 750 to control resistant weeds.

EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with Spraymate Tank & Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

COMPATIBILITY/TANK MIXES

This product can be tank mixed with bromoxynil 200, dicamba, flowable Diuron, methabenzthiazuron, metsulfuron methyl, paraquat/diquat 250, Pelican, Pickout, Salvation, Sethoxydim, Cheminova Sparta 300 SL, and Unleash. The effectiveness of dimethoate, metasystox, omethoate and phosmet may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of non-ionic surfactant is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds.

Drift Warning: DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops and susceptible trees (eg. Kurrajongs, belahs, eucalypts).

Legume Tolerance: The use of MCPA (Cheminova MCPA 750) is preferred to 2,4-D Amine) or Ester in situations where legumes are undersown in the crop.

Note: The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucerne as damage may result.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the product or used container.

RE-ENTRY PERIOD: Hand weeding should not be performed for 7 days after spray application unless workers wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), chemical resistant gloves and footwear.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Leaflet: The method of disposal of the container depends on the container type. Read the 'Storage and Disposal' instructions on the label that is attached to the container.

Non-returnable containers only: Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

100, 110L: Do not remove or tamper with the dry valves or security seal. Do not contaminate the drum with water or any other foreign matter. After each use of the product ensure that the dry valve coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the drum is empty remove the dry valve coupler and return to the point of purchase.

1000L: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes. Will irritate the skin. When opening the container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet available from www.cheminova.com.au

WARNING

This product must be used, handled and stored strictly as directed in accordance with this label, packaging and other reference material ("Directions"). To the extent permitted by law Cheminova Australia Pty Ltd and its related companies will have no liability for any injury, loss or damage arising from a failure to follow the Directions.

APVMA Approval No: 62913/ 61717

Batch No:

Date of Manufacture:

131112-T

Drummaster logo (relevant pack sizes)

**FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111**