

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Rustler[®] 900 WG

Herbicide

ACTIVE CONSTITUENT: 900 g/kg PROPYZAMIDE

GROUP	D	HERBICIDE
-------	---	-----------

For selective control of certain grasses and broad leaf weeds in Canola, Legume Seed Crops and Pastures, Oilseed Poppies, Lettuce and Turf as per the Directions For Use Table.

CHEMINOVA
AUSTRALIA PTY LTD
ABN 59 472 616 773
ACN 110 199 169
12 JULIUS AVE
NORTH RYDE NSW 2113
PH: 1800 624 597
www.cheminova.com.au

NET CONTENTS: 250 g – 25 kg

DIRECTIONS FOR USE

CROPS	WEEDS	STATE	RATE kg/ha	CRITICAL COMMENTS	
BROADACRE CROPS					
Canola	Annual Ryegrass, Barley Grass, Canary Grass, Fescue, Great Brome, Prairie Grass, Rat's Tail Fescue, Silver Grass, Squirrel Tail, Wild Oats (Canola), Winter Grass.	All States	0.56	Incorporate by sowing (IBS) when weeds are at the pre-emergent stage.	
Legume Pastures (including Lucerne, Clover, Medics) grown for Forage, Hay or Seed Production	Bent Grass, Black-Berry Nightshade, Chickweed, English Couch, Patterson's Curse, Perennial Ryegrass, Seedling Sorrel, Seedling Wireweed, Wild Oats, Yorkshire Fog Grass	NSW, ACT, Tas, SA, WA only	0.56 – 0.83	SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July.	
			0.83 – 1.11	WEEDS: Apply when weeds are at post or pre-emergence stages. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.	
Oilseed Poppies	Ryegrass (resistant to herbicides in group A and B)	Tas only	0.56 in 100-200 L water	Apply as foliar application (using suitable boom spray equipment) from 6-leaf stage to crop row cover. Make only one application per crop. DO NOT apply after commencement of running up.	
VEGETABLES					
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed,	All States	2.5 in 330 – 550 L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
Lettuce (Transplanted)	Portulaca, Winter Grass, Rye Grass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles			Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
		Vic only	1.28 in 550 L	Apply on light sandy soils, low in organic matter.	
TURF AND LAWNS					
Common Couch	Winter Grass	All States	0.7 (7 g/100 m ²)	Rate/Bowling Green 37 x 37 m 94 g	Apply during the Autumn and Winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12 – 25 mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Qld Blue Couch and after 4 weeks. DO NOT RESEED FOR 60 DAYS AFTER TREATMENT. Use ONLY on established turf. DO NOT use on bluegrass, ryegrass, fescue or bent grass lawns.
Queensland Blue Couch		Qld only	0.3 (3 g/100 m ²)	47 g	
Buffalo Grass		All States	0.7 (7 g/100 m ²)	94 g	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

- CANOLA:** HARVEST - NOT REQUIRED WHEN USED AS DIRECTED.
GRAZING - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 12 WEEKS AFTER APPLICATION.
- POPPIES:** HARVEST - NOT REQUIRED WHEN USED AS DIRECTED.
GRAZING - DO NOT GRAZE TREATED POPPY CROPS OR CUT FOR STOCKFOOD.
- ALL OTHER CROPS:** DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

RESISTANCE WEED WARNING

GROUP	D	HERBICIDE
-------	---	-----------

Rustler 900 WG Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is a Group D Herbicide. Some naturally occurring weed biotypes resistant to Rustler 900 WG and Group D Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Rustler 900 WG or Group D Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Cheminova Australia Pty Ltd accepts no liability for any losses that may result from the failure of Rustler 900 WG to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or Cheminova Australia Pty Ltd representative.

SPRAY PREPARATION

Rustler 900 WG is a Water Dispersible Granule formulation to be blended with water and applied as a spray. Granule formulations are designed to be added to large volumes of water and dispersed slowly under agitation. When using larger boomsprays and unable to add granules directly to a large volume in the tank via the top of the boomspray, pre-mixing may be necessary as follows:

- Partly (30%) fill the boomspray tank with water.
- Slowly add required amount of Rustler 900 WG for the area to be sprayed into a pre-mixing vessel while agitating well.
- Ensure high water volumes and DO NOT add to the spray tank until the pre-mix is fully dissolved. This may take several mixes to ensure the correct amount is added to the boomspray tank.
- Add the pre-mixed Rustler 900 WG into the boomspray tank with the agitation system operating.
- Fill up to the correct volume with water.
- The mixture must be kept in suspension at all times by continuous agitation.
- Tank-mixtures should be tested for physical compatibility and efficacy before use.

DO NOT mix with MCPA. If tank mixing, Rustler 900 WG must be in suspension before adding other herbicides, insecticides or surfactants.

Wash sprayer thoroughly after use.

EQUIPMENT

Always ensure the sprayer is clean, using a suitable cleaner as directed. It is recommended that for certain turf situations, up to 100L/ha spray volume is required.

RE-ENTRY PERIOD

DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply to, or allow spray to drift onto crops, pastures or non-target – desirable plants, nor onto agricultural land scheduled for cropping or pasture, as injury to the crop or pasture may occur.

PROTECTION OF LIVESTOCK

DO NOT graze sheep, cut for stock feed (including poultry and pigs) any treated areas after application in poppies; any treated areas for 12 weeks after application in canola; any treated areas for 25 days after application in all other crops.

DO NOT FEED TURF/LAWN CLIPPINGS TO POULTRY OR LIVESTOCK.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilizers, insecticides, fungicides and seeds. Shake empty box into spray tank. DO NOT re-use container or dispose of undiluted chemicals on-site. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, State or Territory government regulations. DO NOT burn empty container or product.

SAFETY DIRECTIONS

Harmful if inhaled. Will irritate the eyes. Avoid contact with eyes and skin. DO NOT inhale dust. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves, face shield or goggles and disposable dust face mask covering mouth and nose. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet available from www.Cheminova.com.au

WARNING

This product must be used, handled and stored strictly as directed in accordance with this label, packaging and other reference material ("Directions"). To the extent permitted by law Cheminova Australia Pty Ltd and its related companies will have no liability for any injury, loss or damage arising from a failure to follow the Directions.

APVMA Approval No: 69519/100892

Batch No:

Date of Manufacture:

® Registered trademark of Cheminova

Drummuster logo – relevant pack sizes

150409-T

FOR 24 HOUR SPECIALIST ADVICE IN EMERGENCY ONLY PHONE 1800 033 111
--